

Upper limits on exosatellites around β Pictoris b

Matthew A. Kenworthy¹,¹★ Rico Landman,¹ Andrew Vanderburg¹,^{2,3} Joseph E. Rodriguez¹,⁴ Jayne L. Birkby¹,⁵ Isabella Macias,^{6,7} Darío González Picos,¹ Sydney A. Jenkins¹,⁴ Elina Kleisioti,^{1,8} Tomas Stolker¹ and Ioannis Koutalios¹

¹Leiden Observatory, Leiden University, Postbus 9513, NL-2300 RA Leiden, the Netherlands

²Center for Astrophysics|Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

³Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

⁴Center for Data Intensive and Time Domain Astronomy, Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA

⁵Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

⁶Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

⁷Department of Astronomy, University of Florida, Bryant Space Science Center, Stadium Road, Gainesville, FL 32611, USA

⁸Faculty of Aerospace Engineering, Delft University of Technology, NL-2629 HS Delft, the Netherlands

Accepted 2026 May 29. Received 2026 May 16; in original form 2026 February 13

ABSTRACT

β Pictoris b is one of the closest known directly imaged gas giant exoplanets with an orbit that is almost edge-on to our line of sight, making it an ideal target for radial velocity monitoring to search for massive exomoons. We measure the radial velocity of β Pictoris b over several epochs between October 2024 and March 2025 by using the cross-correlation of a template spectrum with absorption lines in the planet’s atmosphere, giving a mean precision of 160 m s^{-1} . The resultant set of radial velocities is analysed with a periodogram to search for candidate radial velocity (RV) signals indicating a massive exomoon. Although we do not detect an exomoon signal in our data, our detection limits for a single moon are 80 Earth masses at $P = 1 \text{ d}$ and 1 Jupiter at $P = 200 \text{ d}$, comparable to RV exomoon searches around other substellar companions. The RV limit is comparable with the astrometric exomoon limit at a period of 7 d and a mass of $150 M_{\oplus}$, where for longer periods the astrometric searches have lower mass limits. With an additional observing season, the upgraded CRYogenic InfraRed Echelle Spectrograph (CRIRES+) can detect a planet/moon mass ratio of 10^{-3} ($4 M_{\oplus}$) with a period of up to one day, and can detect a Neptune-mass moon at hundreds of Jupiter radii.

Key words: techniques: radial velocities – stars: individual: Beta Pictoris – exoplanets.

1 INTRODUCTION

Moons are seen around all of the gas giant planets in our Solar system, and their final configuration reflects both the composition of the planet and its formation process (e.g. R. Deienno et al. 2011). Such satellites, therefore, must exist around other gas giant planets in the Galaxy – but while there have been some tantalizing candidates detected, none have been confirmed to date. Exomoons may increase the number of Habitable Zones in a planetary system – their temperature may be raised by the flux of energy from their parent planet (R. Heller & R. Barnes 2013) or by tidal heating. Since our current knowledge of moons comes entirely from Solar system satellites, an exomoon discovery could similarly boost our understanding of the diversity that we see in exoplanetary systems (e.g. N. M. Batalha 2014).

To understand the range of possible exomoon properties, it is useful to consider the three primary formation pathways: accretion from a circumplanetary disc (S. R. Kane, N. R. Hinkel & S. N. Raymond 2013), gravitational capture of interplanetary objects

(e.g. suggested for Triton by Neptune; C. B. Agnor & D. P. Hamilton 2006), and collisions between objects. Moons forming from a circumplanetary disc are expected to be relatively small compared to their host planet (masses less than 10^{-5} that of the planet; R. M. Canup & W. R. Ward 2002). While this formation pathway is highly efficient in the Solar system, moons like Triton likely started off as an interplanetary planetesimal and was captured by Neptune some time later (C. B. Agnor & D. P. Hamilton 2006), resulting in Triton’s anomalously large mass ratio with respect to Neptune (2×10^{-4}).

Several methods have been used to search for exomoons, with several candidates proposed; for an overview see the Review in A. Teachey (2024). Gravitational microlensing by a foreground source with proper motion causes achromatic magnification in the resultant light curve of a distant background source. This led to D. P. Bennett et al. (2014) showing the microlensing event MOA-2011-BLG-262 was consistent with a free-floating planet/moon system closer to the Solar system than the background source, although a more distant stellar mass primary/brown dwarf system could produce the same observations. Model degeneracies also led to K.-H. Hwang et al. (2018) identifying the multi-object microlensing event OGLE-2015-BLG-1459L

* E-mail: kenworthy@strw.leidenuniv.nl

as due to one object lensing three background sources and not a foreground star/planet/moon lensing system.

Directly imaging the thermal emission from tidally heated exomoons (THEMs) is a possibility for very close exoplanet systems (M. A. Peters & E. L. Turner 2013) such as the exoplanet Epsilon Eridani b (E. Kleisioti et al. 2023). Tidally locked THEMs can also be detected on high-inclination orbits around their parent planet; any bright spot of thermal emission caused by volcanic activity will result in photometric modulation of the light curve (E. Kleisioti et al. 2024). Volcanic exomoons can also produce an exosphere that is detected as spectral absorption lines of neutral Sodium and Potassium seen before, during and after the transits of hot Jupiters, as seen for WASP-49 b (A. V. Oza et al. 2019).

Exomoons may transit their parent exoplanet, and searches around free floating (M. A. Limbach et al. 2021, 2024; M. J. Wilson et al. 2025) and bound (D. Kipping et al. 2025a; E. K. Pass et al. 2026) self-luminous exoplanets show plausible eclipses, although the variability of the planets themselves can confound this signal.

Searching for exomoons as small as Ganyমে is possible through the impact an exomoon has on the transit of its parent planet across the stellar disc. This can be seen in both transit timing variations and transit duration variations (D. M. Kipping 2009a, b), and were the main methods behind the Hunt for Exomoons with Kepler (D. M. Kipping et al. 2012, 2013; D. Kipping et al. 2022), but to date no compelling candidates have been found around gas giants with orbital periods less than 50 d (D. M. Kipping et al. 2015). Since it is expected that the planets forming in the 5–20 au regime have attendant moons, it is suspected that the process of their migration caused the moons to be scattered away from the system (V. Dobos et al. 2021). The question of whether wider orbit planets host such satellites arises: Two exomoon candidates have been identified through timing variation and possible detection of their transits, Kepler 1708 b-i and Kepler 1625 b-i (A. Teachey & D. M. Kipping 2018; A. Teachey, D. M. Kipping & A. R. Schmitt 2018; D. Kipping et al. 2022, 2025b), around longer-period gas giant exoplanets. These detections are contested: Other teams have carried out an analysis on the data and do not find the exomoon signal (L. Kreidberg, R. Luger & M. Bedell 2019; R. Heller & M. Hippke 2024).

Directly imaged exoplanets offer a greater potential for satellite detection due to their typically much larger separations from the host star, which increases the planet’s Hill sphere radius (V. Dobos et al. 2021), and the absence of significant inward planet migration keeps any attendant moons after their formation. Astrometric searches with optical interferometers can detect exomoons at larger semimajor distances (T. O. Winterhalder et al. 2026), and an initial search around the substellar companion HD 206893 B with the Very Large Telescope Interferometer (VLTI) and GRAVITY+ has resulted in a tantalizing signal (Q. Kral et al. 2026).

Radial velocity (RV) searches have looked at exoplanets in the HR 8799 system (A. Vanderburg & J. E. Rodriguez 2021), HR 7672 B (J.-B. Ruffio et al. 2023), and GQ Lup B (K. Horstman et al. 2024) with no detections. The closest known self-luminous gas giant exoplanet is $\sim 11 M_{\text{Jup}}$ β Pictoris b (A. Lagrange et al. 2009; S. Lacour et al. 2021), located 19 parsecs away around a young (23 Myr; C. P. M. Bell, E. E. Mamajek & T. Naylor 2015) A-type star and embedded in an edge-on circumstellar disc. The orbit of β Pictoris b is highly inclined, with its orbit seen almost close to edge-on (S. Lacour et al. 2021). Building on this geometry, M. Poon, H. Rein & D. Pham (2024) propose that the planet has a large obliquity, given the measured projected rotational velocity

($v \sin i = 20 \text{ km s}^{-1}$; R. Landman et al. 2024), and suggest that a Neptune-mass exomoon at 40 to 70 planetary radii could be the cause. RV measurements measure the mass function $M \sin i$ and for exomoons we have no measurements for any orbital inclinations. Moons in the Solar system are generally coplanar with the equator of their parent planet, while β Pictoris b has a relatively high inclination as measured from the rotational broadening of absorption lines within its atmosphere (I. A. G. Snellen et al. 2014), so it is not an unreasonable assumption to assume any moons around the planet will also have a high inclination as seen from Earth.

In 2016, the planet’s Hill sphere transited the star to within 20 per cent of the Hill sphere radius, and a photometric monitoring programme established that there was no significant circumplanetary dust (M. A. Kenworthy et al. 2021), concluding that any material had already condensed into satellites around the planet and may well have high inclinations amenable for detection with RV methods.

We present upper limits on our search for exomoons around β Pictoris b using the upgraded CRYogenic InfraRed Echelle Spectrograph (CRIRES+) combined with astrometric limits derived from archival astrometric data (I. Macias, S. A. Jenkins & A. Vanderburg 2026). In Section 2 we detail the observations with the VLT, followed by a periodogram analysis with injection and recovery to determine sensitivity limits. We discuss these results in Section 3 along with our conclusions in Section 4.

2 OBSERVATIONS AND ANALYSIS

β Pictoris b was observed as a service mode programme with various epochs between UT 2024 Oct 17 and UT 2025 Mar 09 using VLT/CRIRES+. We used the $0.2''$ slit and the K2166 wavelength setting. The slit was oriented on the planet location at a 45° deg angle with respect to the line connecting the star and the planet. This allows us to avoid overexposure from the bright star, while still allowing us to use the starlight for calibration purposes. Each observation consisted of 24 integrations of 120 s in an AAABBBBBBAAA nodding scheme. We chose this approach to minimize the overhead time from the nodding cycles, while still allowing for background subtraction, which is done using the average of the three closest exposures in the other nodding position. We analyse the exposures from the A and B frames separately, as they can have slightly different wavelength solutions.

We reduced the spectra with two separate pipelines. In the first one, the initial data reduction, such as background subtraction, flat fielding, and a first wavelength solution, was done with *pycrises* (T. Stalker & R. Landman 2023), which makes use of the official CRIRES+ pipeline.¹ We follow the reduction and analysis steps detailed in R. Landman et al. (2024) to detect the planet and obtain an estimate of its radial velocity, and give a short summary here: First, we refine the wavelength solution using the telluric lines imprinted on the spectra of the host star. The stellar contamination at the location of the planet is estimated using low-pass filtering and a Principal Component Analysis (PCA). We use 10 PCA components to model the stellar and telluric contribution. We then jointly fit the stellar contribution and planet signal using the likelihood framework developed in J.-B. Ruffio et al. (2021) and adapted to CRIRES+ in R. Landman et al. (2024). In this

¹<https://www.eso.org/sci/software/pipelines/crises/crises-pipe-recipes.html>




Figure 1. A section of the spectrum of β Pictoris b from CRIRES+, showing one night of data and the fitted model. The template spectra of water and carbon monoxide are shown above, indicating their relative contributions.

case, we fix all the parameters of the planet, e.g. the atmospheric parameters and rotational velocity, to the values obtained in R. Landman et al. (2024). This likelihood was calculated for a grid of radial velocity values, which was finally converted to a maximum likelihood estimate and an associated uncertainty. The mean precision was 320 m s^{-1} .

A second, separate and fully independent reduction and analysis were carried out to validate the RV measurements. In this workflow, the data were processed using the Python package EXCALIBUR (Y. Zhang et al. 2024). The reduction steps closely follow those of *pycrites*; however, we implemented a revised extraction scheme that combines all spatial pixels containing a significant planetary signal. A section of one of the reduced spectra is shown in Fig. 1. Atmospheric retrievals were then performed using forward models generated with *petitRADTRANS* (P. Mollière et al. 2019) and explored via nested sampling with *ultranest* (J. Buchner 2021). This approach allows us to constrain the planet’s atmospheric properties, radial velocity, and rotational broadening. We adopted state-of-the-art line lists for water (H_2O ; O. L. Polyansky et al. 2018) and carbon monoxide (CO ; L. S. Rothman et al. 2010; G. Li et al. 2015). The atmospheric composition is parametrized using constant-with-altitude abundances, with free parameters for each relevant molecular species, as described in S. de Regt et al. (2024). The temperature structure is modelled using a parametrized profile in which temperature gradients at reference atmospheric layers are retrieved, following the implementation of D. González Picos et al. (2024). The complete results and interpretation of the atmospheric retrievals, together with details of the improved spectral extraction scheme, will be provided in a forthcoming paper (Gonzalez Picos et al., accepted). The contribution of starlight at the planet’s location is incorporated directly into the forward model, following the approach demonstrated for GQ Lup B by D. González Picos et al. (2025), adapted from J.-B. Ruffio et al. (2023) and R. Landman et al. (2024). We adopt these radial velocities for subsequent analysis due to their greater precision.

In Fig. 2, we show the detection of the planet’s rotation and the identification of water (H_2O) and carbon monoxide (CO), which are the dominant opacity sources over the wavelength range of the observations ($\lambda = 2060\text{--}2472 \text{ nm}$). The planetary spectrum and the best-fitting model are shown in Fig. 1. For subsequent analysis, we adopt these physical parameters of β Pictoris b: from S. Lacour et al. (2021) we take the mass of β Pictoris b to be $11.9 \pm 3.0 M_{\text{Jup}}$, with orbital parameters $a = 9.93 \pm 0.03 \text{ au}$, $e = 0.103 \pm 0.003$, and $i = 89.00 \pm 0.01 \text{ deg}$. For the radius, we adopt the Exo-REM posterior from M. Ravet et al. (2025) of 1.73 ± 0.01




Figure 2. Cross-correlation of the CRIRES+ spectrum of the planet, (a) indicating the rotational broadening of the planet and at the retrieved $v \sin i \approx 19 \text{ km s}^{-1}$ and the autocorrelation function of the unbroadened model spectrum. (b) The relative detection strengths of the planet in water and carbon monoxide.

R_{Jup} . We calculate the Roche limit both for solid and fluid bodies using the mass/radius relationship from J. Chen & D. Kipping (2017), although we note that young objects may have a larger radius and therefore lower density.

Analysis of the RV measurements (see Fig. 3) was done using the RVSEARCH Python package (L. J. Rosenthal et al. 2021). The algorithm fits both a ‘no-moon’ model and ‘one moon’ model for




Figure 3. Radial velocity measurements of β Pictoris b from CRIRES+ with the D. González Picos et al. (2024) formulation.




Figure 4. Results of the blind search for exosatellites in the RV time series listed in Table 1 using RVSEARCH. It shows the difference in the Δ BIC between a model including a satellite and a model without a satellite. We also show the threshold for the 0.001 false alarm probability (FAP).

a given set of trial periods from 1 to 500 d in a circular orbit, and the difference between these two Bayesian information criterion (BIC) models, Δ BIC, is calculated. It then estimates a Δ BIC detection threshold that corresponds to a false alarm probability of 0.1 per cent, following the methodology outlined in A. W. Howard & B. J. Fulton (2016) where further details can be found. The results are shown in Fig. 4. Even though the shortest time between two observations is around two days, we are sensitive to shorter orbital periods due to the irregularly spaced sampling of our measurements. The largest Δ BIC has a value of 12 for an orbital period of 2.3 d, below the false alarm probability threshold of 46. A set of injection and recovery tests for various trial periods and masses for a single exomoon on a circular orbit was performed, with the resultant recovery limits shown in Fig. A1. Fig. 5 shows the upper limits for a single exomoon orbiting β Pictoris b in a circular orbit, as a function of semimajor axis and moon to planet mass function ratio.

3 DISCUSSION

The mass ratio of moons to planets is around 10^{-4} for the gas giants (R. M. Canup & W. R. Ward 2006), but theoretical studies suggest that more massive planets have more massive moons with a scaling law of $M_{\text{moon}} \propto M_p^{3/2}$ with ratios approaching 10^{-3} (see

equation 43 in K. Batygin & A. Morbidelli 2020). From the RV analysis, no compelling periodic signal is seen in the data presented in the previous section. The smallest mass ratio reached is 2×10^{-2} ($60 M_{\oplus}$) at the fluid Roche limit, just below a Saturn mass exomoon with an orbital period of 0.9 d, gradually increasing to a mass ratio of just under 0.1 at 250 d. The sensitivity is comparable to previous RV searches around other substellar and planetary companions. A study on the HR 8799 b, c, d exoplanets with Keck OSIRIS radial velocities ruled out most companions with $m \sin i > 2 M_{\text{Jup}}$ for orbital periods shorter than 5 d (A. Vanderburg & J. E. Rodriguez 2021) whereas we reach down to just over a Saturn mass for β Pictoris b, whilst a limit of 10^{-2} ($100 M_{\oplus}$) was reached for orbital periods of 1 d around the substellar companion GQ Lup B (K. Horstman et al. 2024).

An additional observing campaign, consisting of 25 CRIRES+ observations sampled randomly over a 50 d period, was simulated and added to the current RV data set, and the resultant sensitivity curve is shown in Fig. 5 with the injection recovery limits shown in Fig. A2. Given the previous performance of CRIRES+, we can reach a mass ratio of 10^{-3} for an orbital period of 1 d, and we can potentially detect Neptune mass exomoons out to periods of 200 d.

I. Macias et al. (2026) analysed archival astrometric data on β Pictoris b and determined upper limits on exomoons. These limits are shown in Fig. 5. Astrometric detection limits complement the radial velocity methods, and we can now rule out $150 M_{\oplus}$ exomoons for edge-on circular orbits around β Pictoris b for all prograde stable orbits (estimated as 0.44 of the Hill sphere radius using equation 5 from R. C. Domingos, O. C. Winter & T. Yokoyama 2006). Indeed, even though there is a peak in the periodogram at around 52 d (but still below the FAP limit), the astrometric limits are far more constraining at this period and show no signal.

4 CONCLUSIONS

These reported measurements represent the most sensitive limits to exomoons at short orbital periods to date for β Pictoris b, and the complementarity of astrometric limits give coverage throughout the Hill sphere of the planet. Extending the monitoring of β Pictoris b with CRIRES+ will increase the sensitivity to moons with mass ratios greater than about 5×10^{-3} i.e. moons that formed/arrived via gravitational capture pathways or ones which induce the obliquity in β Pictoris b (M. Poon et al. 2024) – see Fig. 5. Together with the first astrometric limits of β Pictoris b from I. Macias et al. (2026) we can rule out any exomoons three times more massive than Saturn in the Hill sphere.

The discovery of massive gas giant exoplanets at large separations from their parent stars, such as HD106096 b (V. Bailey et al. 2014), YSES-1 b and c (A. J. Bohn et al. 2020), and WISPIIT 1 b and c (R. F. van Capelleveen et al. 2025) enable direct spectroscopic monitoring of the planets to look for exosatellites. One highly promising candidate is YSES 1b which hosts both a circumplanetary disc seen in thermal emission (K. K. W. Hoch et al. 2025) and whose orbit is almost edge on to our line of sight (J. Roberts et al. 2025). If the geometry is particularly favourable and the moon transits the disc of the exoplanet, the Rossiter–McLaughlin effect can produce a very pronounced RV signal due to the rapid rotation of these young exoplanets (J.-B. Ruffio et al. 2023). If exomoons do transit β Pictoris b and other wide projected separation planets such as YSES 1 b, then we may be highly fortunate and have our first RV-detected exomoons within the next few years.




Figure 5. The upper limits on exosatellites around β Pictoris b. The blue line represents the radial velocity limits for 50 per cent retrieval of injected edge-on circular orbits. The red line is the outermost stable prograde orbit for an exosatellite around β Pictoris b. The pale green line corresponds to astrometric limits from I. Macias et al. (2026). The Roche limits for rigid and fluid exosatellite bodies are shown in grey on the left. The predicted mass and location for an exomoon which could give Beta Pic b a high obliquity is indicated in purple (M. Poon et al. 2024). The predicted sensitivity for an additional 25 observations with CRILES+ is shown in purple.

Table 1. Radial velocity measurements of β Pictoris b.

MJD	RV _{instrument} (km s ⁻¹)	RV _{corr} (km s ⁻¹)	RV _{helio} (km s ⁻¹)
60600.36	26.00 ^{+0.11} _{-0.10}	+6.78	32.78
60604.35	25.81 ^{+0.07} _{-0.07}	+6.46	32.27
60683.10	35.17 ^{+0.09} _{-0.08}	-3.39	31.78
60686.23	36.11 ^{+0.11} _{-0.13}	-4.00	32.11
60702.20	38.46 ^{+0.13} _{-0.16}	-5.84	32.66
60733.07	39.13 ^{+0.08} _{-0.08}	-7.92	31.20
60735.12	39.83 ^{+0.13} _{-0.13}	-8.06	31.77
60739.08	40.13 ^{+0.15} _{-0.15}	-8.12	32.01
60743.08	40.51 ^{+0.14} _{-0.16}	-8.20	32.31
60755.06	39.97 ^{+0.08} _{-0.11}	-8.16	31.81
60760.01	39.41 ^{+0.10} _{-0.09}	-7.99	31.42

ACKNOWLEDGEMENTS

We thank our referee for their careful reading of our manuscript and suggestions which improved it. JLB acknowledges funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme under grant agreement no. 805445, and the support of the Leverhulme Trust via the Philip Leverhulme Physics Prize. Based on observations collected at the European Southern Observatory under ESO programme 114.27DX.001. This research has used the SIMBAD data base, operated at CDS, Strasbourg, France (M. Wenger et al. 2000). This work has used data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consor-

tium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. To achieve the scientific results presented in this article, we made use of the *Python* programming language,² especially the *SCIPY* (P. Virtanen et al. 2020), *NUMPY* (T. E. Oliphant 2007), *MATPLOTLIB* (J. D. Hunter 2007), *EMCEE* (D. Foreman-Mackey et al. 2013), *ASTROPY* (Astropy Collaboration 2013; A. M. Price-Whelan et al. 2018) packages.

DATA AVAILABILITY

This paper is compiled using the *showyourwork!* framework, which includes all original data files, computer scripts to generate the figures, and associated compilation framework.

This is available at <https://github.com/mkenworthy/BetaPicbExomoons>

REFERENCES

- Agnor C. B., Hamilton D. P., 2006, *Nature*, 441, 192
- Astropy Collaboration, 2013, *A&A*, 558, A33
- Bailey V. et al., 2014, *ApJ*, 780, L4
- Batalha N. M., 2014, *Proc. Natl. Acad. Sci. USA*, 111, 12647
- Batygin K., Morbidelli A., 2020, *ApJ*, 894, 143
- Bell C. P. M., Mamajek E. E., Naylor T., 2015, *MNRAS*, 454, 593
- Bennett D. P. et al., 2014, *ApJ*, 785, 155
- Bohn A. J. et al., 2020, *ApJ*, 898, L16
- Buchner J., 2021, *J. Open Source Softw.*, 6, 3001
- Canup R. M., Ward W. R., 2002, *AJ*, 124, 3404
- Canup R. M., Ward W. R., 2006, *Nature*, 441, 834

²Python Software Foundation, <https://www.python.org/>

Chen J., Kipping D., 2017, *ApJ*, 834, 17
 de Regt S. et al., 2024, *A&A*, 688, A116
 Deienno R., Yokoyama T., Nogueira E. C., Callegari N., Santos M. T., 2011, *A&A*, 536, A57
 Dobos V., Charnoz S., Pál A., Roque-Bernard A., Szabó G. M., 2021, *PASP*, 133, 094401
 Domingos R. C., Winter O. C., Yokoyama T., 2006, *MNRAS*, 373, 1227
 Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
 González Picos D. et al., 2024, *A&A*, 689, A212
 González Picos D., Snellen I. A. G., de Regt S., Landman R., Zhang Y., Gandhi S., Sánchez-López A., 2025, *A&A*, 693, A298
 Heller R., Barnes R., 2013, *Astrobiology*, 13, 18
 Heller R., Hipke M., 2024, *Nat. Astron.*, 8, 193
 Hoch K. K. W. et al., 2025, *Nature*, 643, 938
 Horstman K. et al., 2024, *AJ*, 168, 175
 Howard A. W., Fulton B. J., 2016, *PASP*, 128, 114401
 Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
 Hwang K.-H. et al., 2018, *AJ*, 155, 259
 Kane S. R., Hinkel N. R., Raymond S. N., 2013, *AJ*, 146, 122
 Kenworthy M. A. et al., 2021, *A&A*, 648, A15
 Kipping D. M., 2009a, *MNRAS*, 392, 181
 Kipping D. M., 2009b, *MNRAS*, 396, 1797
 Kipping D. M., Bakos G. Á., Buchhave L., Nesvorný D., Schmitt A., 2012, *ApJ*, 750, 115
 Kipping D. M., Hartman J., Buchhave L. A., Schmitt A. R., Bakos G. Á., Nesvorný D., 2013, *ApJ*, 770, 101
 Kipping D. M., Schmitt A. R., Huang X., Torres G., Nesvorný D., Buchhave L. A., Hartman J., Bakos G. Á., 2015, *ApJ*, 813, 14
 Kipping D., Cassese B., Changeat Q., Yahalom D., Teachey A., Edwards B., 2025a, preprint (arXiv:2511.15317)
 Kipping D. et al., 2022, *Nat. Astron.*, 6, 367
 Kipping D. et al., 2025b, *Nat. Astron.*, 9, 795
 Kleisioti E., Dirx D., Rovira-Navarro M., Kenworthy M. A., 2023, *A&A*, 675, A57
 Kleisioti E., Dirx D., Tan X., Kenworthy M. A., 2024, *A&A*, 687, A125
 Kral Q. et al., 2026, *A&A*, 705, A217
 Kreidberg L., Luger R., Bedell M., 2019, *ApJ*, 877, L15
 Lacour S. et al., 2021, *A&A*, 654, L2
 Lagrange A. et al., 2009, *A&A*, 506, 927
 Landman R. et al., 2024, *A&A*, 682, A48
 Li G., Gordon I. E., Rothman L. S., Tan Y., Hu S.-M., Kassi S., Campargue A., Medvedev E. S., 2015, *ApJS*, 216, 15
 Limbach M. A., Vos J. M., Vanderburg A., Dai F., 2024, *AJ*, 168, 54
 Limbach M. A., Vos J. M., Winn J. N., Heller R., Mason J. C., Schneider A. C., Dai F., 2021, *ApJ*, 918, L25
 Macías I., Jenkins S. A., Vanderburg A., 2026, *AJ*, 171, 197
 Mollière P., Wardenier J. P., van Boekel R., Henning T., Molaverdikhani K., Snellen I. A. G., 2019, *A&A*, 627, A67
 Oliphant T. E., 2007, *Comput. Sci. Eng.*, 9, 10
 Oza A. V. et al., 2019, *ApJ*, 885, 168
 Pass E. K., Charbonneau D., Vanderburg A., Bean J. L., 2026, preprint (arXiv:2604.05235)
 Peters M. A., Turner E. L., 2013, *ApJ*, 769, 98
 Polyansky O. L., Kyuberis A. A., Zobov N. F., Tennyson J., Yurchenko S. N., Lodi L., 2018, *MNRAS*, 480, 2597
 Poon M., Rein H., Pham D., 2024, *Open J. Astrophys.*, 7, 109
 Price-Whelan A. M. et al., 2018, *AJ*, 156, 123
 Ravet M. et al., 2025, *A&A*, 704, A325
 Roberts J. et al., 2025, *AJ*, 170, 273
 Rosenthal L. J. et al., 2021, *ApJS*, 255, 8
 Rothman L. S. et al., 2010, *J. Quant. Spec. Radiat. Transf.*, 111, 2139
 Ruffio J.-B. et al., 2021, *AJ*, 162, 290
 Ruffio J.-B. et al., 2023, *AJ*, 165, 113
 Snellen I. A. G., Brandl B. R., de Kok R. J., Brogi M., Birkby J., Schwarz H., 2014, *Nature*, 509, 63
 Stolk T., Landman R., 2023, *Astrophysics Source Code Library*, preprint (ascl:2307.040)
 Published by Oxford University Press on behalf of Royal Astronomical Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Teachey A., 2024, preprint (arXiv:2401.13293)
 Teachey A., Kipping D. M., 2018, *Sci. Adv.*, 4, eaav1784
 Teachey A., Kipping D. M., Schmitt A. R., 2018, *AJ*, 155, 36
 van Capelleveen R. F. et al., 2025, *A&A*, 704, A221
 Vanderburg A., Rodriguez J. E., 2021, *ApJ*, 922, L2
 Virtanen P. et al., 2020, *Nat. Methods*, 17, 261
 Wenger M. et al., 2000, *A&AS*, 143, 9
 Wilson M. J. et al., 2025, *AJ*, 170, 357
 Winterhalder T. O. et al., 2026, *A&A*, 705, A216
 Zhang Y. et al., 2024, *AJ*, 168, 246

APPENDIX A: INJECTION RECOVERY TESTS




Figure A1. Exosatellite detection limits derived using injection and recovery tests for β Pictoris b.




Figure A2. Exosatellite detection limits with 25 additional epochs of CRIRES+ observations with precision of 250 m s^{-1} in the 2025/26 observing season.

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.

© The Author(s) 2026.