

OAI 2015

Exercises on Slit Spectrographs

M.A. Kenworthy

11 November 2015

1. The MMTO telescope has a diameter of 6.5m and a Cassegrain focal ratio of $f/15$. A spectrograph at the Cassegrain focus has an entrance slit set to match the typical seeing of 0.75 arcseconds.
 - (a) How wide is the slit in mm? [1]
 - (b) The diameter of the collimator is 300mm, and it projects onto a glass prism made of F2 glass with a base length $B = 200\text{mm}$. What is the dispersion A in radians per nm at the Hydrogen alpha line? You can assume $dn/dl = 0.06\mu\text{m}^{-1}$ at a wavelength of 0.656 microns and the prism is set up in minimum deviation. [2]
 - (c) The camera for the spectrograph is at $f/2$. What is the slit width on the detector? [2]
 - (d) The detector is a CCD with 512 pixels to a side and 20 micron pixel size. Is the CCD adequately sampling the slit? [2]
 - (e) What is the spectral resolution of the spectrograph at 656nm? [2]
 - (f) What is the estimated bandwidth of the spectrograph? [1]
 - (g) Why can you only estimate the bandwidth? [1]
2. The James Webb Space Telescope has a diameter of 6.5m and a focal length of 130m.
 - (a) What is the size of the Airy disk at the telescope focus for a wavelength of 2.2 microns? [1]
 - (b) What is the width of the spectrograph entrance slit that matches this diffraction limit? [2]
 - (c) Studying the dynamics of globular clusters requires a spectral resolution of 60000. A diffraction grating with 100 lines/mm is provided for the spectrograph. If the diffraction grating is used in 8th order in a Littrow configuration, what is the incident angle on the grating? [2]
 - (d) What is the number of illuminated grooves needed for this spectral resolution? [2]
 - (e) What is the full length of the grating? [1]
 - (f) What is the diameter of the collimated beam and its focal length? [2]