

Hand in: Thursday 2 October

Homework exercises OAF2

Bose Einstein statistics

Exercise 3

- a) Show $X(t)$ that the probability density function for the intensity of an unpolarised signal from a thermal source is given by:

$$p(I) = \bar{I}^{-1} e^{-I/\bar{I}}, \quad (1)$$

where $\bar{I} = 2\sigma^2$. The variance σ^2 here relates to the (uncorrelated) gaussian stochastic variables $\text{Re}(\tilde{E}_0(t))$ and $\text{Im}(\tilde{E}_0(t))$, that give a scalar description of the fluctuating behaviour of the *phasor* $\tilde{E}_0(t)$.

- b) Proof that for the variance of the intensity I of an unpolarised signal:

$$\overline{\Delta I^2} = (\bar{I})^2 \quad (2)$$

Exercise 4

- a) The equation for the stationary state of a free particle follows from the solution of the time-independent Schrödinger equation. The wave amplitude Ψ is then given by:

$$\Psi(\vec{r}) = e^{\frac{2\pi i}{h} \vec{p} \cdot \vec{r}} \quad (3)$$

When considering a finite volume, the allowed values for the momentum \vec{p} must meet the specific boundary conditions, which causes quantization of the allowed momentum and energy levels.

Show that the number of occupation levels dZ_k for values of the momentum between p_k and $p_k + dp_k$ *per unit of volume* is given by:

$$dZ_k = \frac{4\pi p_k^2}{h^3} dp_k \quad (4)$$

- b) Consider a black body cavity with temperature T . The radiation field can be described as a *boson gas*. Calculate the number of occupation levels per cm^3 for the *average* photon frequency that corresponds to a temperature of $T = 5000K$, integrated over the frequency interval of the natural line width. Assume a coherence time of $\Delta t_c = 10^{-8}$ s.
- c) Derive the *energy density per Hz* of the black body radiation field, using the result you found above and the mean occupation number for a photon gas.