

Christoph Keller, C.U.Keller@astro.uu.nl

- [www.astro.uu.nl/~keller](http://www.astro.uu.nl/~keller)
- Solar Physics
  - observations of solar magnetic fields
  - analysis of realistic MHD simulations
- Instrumentation
  - Dutch Open Telescope (DOT) at La Palma, Canary Islands
  - polarimeters for solar magnetic field studies
  - direct imaging of exoplanetary systems with large telescopes
  - high-resolution techniques such as Adaptive Optics

## Literature

- Michael Stix, **The Sun: An Introduction**, Second Edition, Springer
- PDF file of this lecture with links at [www.astro.uu.nl/~keller/Teaching/USAP2006/TheSun.pdf](http://www.astro.uu.nl/~keller/Teaching/USAP2006/TheSun.pdf)

# Our Sun: a Star Close-Up

## Outline

- **The Sun: An Introduction (9:30-10:15)**
  - The Sun's Uniqueness
  - Solar Structure and Terminology
  - Current Problems in Solar Physics
- **The Sun as a Star (10:30–11:15)**
  - The Sun among the Stars
  - Evolution of the Sun
  - Solar Neutrinos
- **The Sun: A Magnetic Star (11:30-12:15)**
  - Flux Tube Observations and Theory
  - Faculae
  - The Solar Cycle
- **Solar Telescope Design (14:00–17:00)**

# What makes the Sun Unique?


# What makes the Sun Unique?

## Some Answers

- Sun is the closest star
- Only star with well-resolved atmosphere
  - electromagnetic radiation
  - particle detection
- Only star with well-observed interior
  - helioseismology
  - neutrinos
- Only star of importance for life on Earth




# Solar Structure and Terminology




## Basic Facts

Solar radius	695,990 km	109 Earth radii
Solar mass	$1.989 \cdot 10^{30}$ kg	333,000 Earth masses
Solar luminosity	$3.846 \cdot 10^{33}$ erg/s	
Surface temperature	5770 K	
Surface density	$2.07 \cdot 10^{-7}$ g/cm <sup>3</sup>	$1.6 \cdot 10^{-4}$ Air density
Surface composition	70% H, 28% He, 2% CNO by mass	
Central temperature	15,600,000 K	
Central density	150 g/cm <sup>3</sup>	8 times Gold density
Central composition	35% H, 63% He, 2% CNO by mass	
Solar age	$4.57 \cdot 10^9$ yr	


## The Photosphere



# The Chromosphere



## The Corona seen during a Solar Eclipse




[antwrp.gsfc.nasa.gov/apod/ap060407.html](http://antwrp.gsfc.nasa.gov/apod/ap060407.html)

## The Corona in 1992 seen in X-Rays from the Yohkho Satellite




[www.windows.ucar.edu/cgi-bin/tour\\_def/sun/atmosphere/corona.html](http://www.windows.ucar.edu/cgi-bin/tour_def/sun/atmosphere/corona.html)

# The Solar Wind




## Sunspots






## Flares



[sohowww.nascom.nasa.gov/hotshots/2003\\_10\\_28/](http://sohowww.nascom.nasa.gov/hotshots/2003_10_28/)


## Coronal Mass Ejection




2003/10/28 00:18

[sohowww.nascom.nasa.gov/hotshots/2003\\_10\\_28/](http://sohowww.nascom.nasa.gov/hotshots/2003_10_28/)

# Rotation



[galileo.rice.edu/sci/observations/sunspot\\_drawings.html](http://galileo.rice.edu/sci/observations/sunspot_drawings.html)



[soi.stanford.edu/press/GONG\\_MDI\\_03-00/](http://soi.stanford.edu/press/GONG_MDI_03-00/)

## Differential Rotation


- Christoph Scheiner in 1630: slower rotation at higher latitudes
- helioseismology reveals internal solar rotation rate
- only convection zone shows differential rotation

## Current Problems in Solar Physics

- **oxygen abundance:** numerical simulations imply metal abundances that are in disagreement with helioseismic frequencies
- **FIP-effect:** photospheric and solar wind abundances are not the same
- **origin of supergranulation:** physical mechanism
- **coronal heating process:** energy source, transport, dissipation mechanisms
- **solar wind acceleration:** physical mechanism
- **Nature of Flares:** source of magnetic energy, instability, forecasting
- **origin of solar cycle:** physics of the (large-scale) dynamo
- **origin of small-scale fields:** leftovers from sunspot cycle or small-scale dynamo in surface layers

# The Sun among the Stars


## An Artist's Impression



[trace.lmsal.com/POD/NAS2002\\_otherimages.html](http://trace.lmsal.com/POD/NAS2002_otherimages.html)

# Magnetic Activity

## The 11-Year Solar Cycle





[trace.lmsal.com/POD/NAS2002\\_otherimages.html](http://trace.lmsal.com/POD/NAS2002_otherimages.html)

*If the Sun had no magnetic field,  
it would be as uninteresting as many astronomers think it is.*

R.B.Leighton: unpublished remark (ca. 1965)

## Irradiance and Sunspots







[earthobservatory.nasa.gov/Newsroom/NasaNews/2003/2003032011367.html](http://earthobservatory.nasa.gov/Newsroom/NasaNews/2003/2003032011367.html)


## The Solar (not so) Constant

- correlation between irradiance and magnetic activity
- sunspots only temporarily reduce irradiance
- faculae more than compensate sunspot deficit
- solar constant varies by about 0.1%

# Stellar Irradiance vs. Magnetic Variations




## Doppler Imaging



[www.astro.uu.se/~oleg/structures.html](http://www.astro.uu.se/~oleg/structures.html)

- quickly rotating stars
- many spectra per rotation period
- fit with 'spotted' star model
- also possible for polarized spectra

## Zeeman Doppler Imaging




[www.ast.obs-mip.fr/users/donati/map.html](http://www.ast.obs-mip.fr/users/donati/map.html)



# Evolution of the Sun

## The Past Sun

- numerical models of stellar evolution
  - include all relevant physics including rotation
  - solve PDEs for each time step
  - adjust abundances after each time step
  - have to produce currently observed Sun
- current age:  $4.57 \times 10^9$  years (22 times around galaxy center)
- very young Sun:
  - 70% of current luminosity
  - 125 K colder surface
  - 13% smaller
  - very active chromosphere and corona
  - strong solar wind
  - rotation period only 9 days
- how could life on Earth start and survive?



[www.astro.uva.nl/demo/sun/leven.htm](http://www.astro.uva.nl/demo/sun/leven.htm)

## The Future of the Sun

- B** 50% of available H in core used up (now)
- C** all H in core used up, H fusion in shell, 40% larger, twice as bright
- D**  $1.5 \times 10^9$  years later 3 times normal size, temperature 4300 degrees;  $0.25 \times 10^9$  years later, 100 times larger, 500 times more luminous
- E** critical core temperature, all He fuses into C, explosion throws out  $\frac{1}{3}$  of solar mass into space, planetary nebula and white dwarf ( $2000 \text{ kg/cm}^3$ )


## Solar Neutrinos

- no direct electromagnetic radiation from solar interior
- luminosity, age of the Sun ( $4.5 \times 10^9$  years)  $\Rightarrow$  nuclear fusion
- knowledge of stellar interiors largely based on model calculations
- *helioseismology* measures sound speed as function of depth
- neutrinos from solar core are the only direct measurement
- $\nu_e$  are produced in fusion reactions, leave the Sun within 2.3 s
- on Earth each  $\text{cm}^2$  is penetrated every second by  $4 \cdot 10^{10}$  solar  $\nu_e$
- $\nu$ -telescope makes  $\nu$ -images of the solar core
- weak interaction only,  $\nu$  hard to detect  $\Rightarrow$  very large detectors

# The Problem

## The Data

Total Rates: Standard Model vs. Experiment  
Bahcall-Serenelli 2005 [BS05(OP)]



[www.sns.ias.edu/jnb/SNviewgraphs](http://www.sns.ias.edu/jnb/SNviewgraphs)

## Observations and Standard Models Disagree


- Wrong standard solar model? But helioseismology excludes non-standard solar core models
- Wrong standard model of particle physics? But neutrinos might have mass

## The Solution


- Mikheyev–Smirnov–Wolfenstein effect
- interaction with matter:  $\nu_e$  and  $e^-$  can interact through  $W^-$  or  $Z^0$ ,  $\nu_\mu$  and  $\nu_\tau$  can only interact with  $e^-$  through  $Z^0$
- most neutrinos from Sun will pass through resonance density region inside the Sun
- even very small mixing angles and mass differences can make most  $\nu_e$  into  $\nu_\mu$
- all solar data and also reactor experiments deliver consistent combinations of mixing angle and difference of squared masses

# Flux Tubes: Observational Evidence


## DOT Call K image close to the limb



# DOT H $\alpha$ image




## TRACE Loops






## SOLIS VSM Magnetic Field Distribution




## Direct Detection of Concentrated Fields



# Evidence from MHD Simulations

## Stein & Nordlund, Quiet Sun



# Thin Flux Tube Approximation

## Force Balance

- all relative length scales are large compared to tube diameter
- neglect diffusion term in induction equation
- equation of motion

$$\rho \left( \frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} \right) = -\nabla p + \vec{j} \times \vec{B} + \vec{F}_{\text{gravity}} + \vec{F}_{\text{viscosity}}$$

- magnetohydrostatic ( $\vec{v} = 0 \Rightarrow F_{\text{viscosity}} = 0$ )
- force balance

$$\nabla p - \vec{j} \times \vec{B} = \vec{F}_{\text{gravity}}$$

- with  $\mu_0 \vec{j} = \nabla \times \vec{B}$  and  $\vec{B} \times (\nabla \times \vec{B}) = \frac{1}{2} \nabla B^2 - (\vec{B} \cdot \nabla) \vec{B}$

$$\nabla \left( p + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} (\vec{B} \cdot \nabla) \vec{B} = \vec{F}_{\text{gravity}}$$

# Thin Flux Tube Approximation

## Force Balance

- all relative length scales are large compared to tube diameter
- neglect diffusion term in induction equation
- equation of motion

$$\rho \left( \frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} \right) = -\nabla p + \vec{j} \times \vec{B} + \vec{F}_{\text{gravity}} + \vec{F}_{\text{viscosity}}$$

- magnetohydrostatic ( $\vec{v} = 0 \Rightarrow F_{\text{viscosity}} = 0$ )
- force balance

$$\nabla p - \vec{j} \times \vec{B} = \vec{F}_{\text{gravity}}$$

- with  $\mu_0 \vec{j} = \nabla \times \vec{B}$  and  $\vec{B} \times (\nabla \times \vec{B}) = \frac{1}{2} \nabla B^2 - (\vec{B} \cdot \nabla) \vec{B}$

$$\nabla \left( p + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} (\vec{B} \cdot \nabla) \vec{B} = \vec{F}_{\text{gravity}}$$

# Thin Flux Tube Approximation

## Force Balance

- all relative length scales are large compared to tube diameter
- neglect diffusion term in induction equation
- equation of motion

$$\rho \left( \frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} \right) = -\nabla p + \vec{j} \times \vec{B} + \vec{F}_{\text{gravity}} + \vec{F}_{\text{viscosity}}$$

- magnetohydrostatic ( $\vec{v} = 0 \Rightarrow F_{\text{viscosity}} = 0$ )
- force balance

$$\nabla p - \vec{j} \times \vec{B} = \vec{F}_{\text{gravity}}$$

- with  $\mu_0 \vec{j} = \nabla \times \vec{B}$  and  $\vec{B} \times (\nabla \times \vec{B}) = \frac{1}{2} \nabla B^2 - (\vec{B} \cdot \nabla) \vec{B}$

$$\nabla \left( p + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} (\vec{B} \cdot \nabla) \vec{B} = \vec{F}_{\text{gravity}}$$

## Radial Force Balance

- force balance in general coordinate system

$$\nabla \left( p + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} \left( \vec{B} \cdot \nabla \right) \vec{B} = \vec{F}_{\text{gravity}}$$

- in cylindrical coordinates, radial component

$$\frac{\partial}{\partial r} \left( p + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} \left( B_r \frac{\partial B_r}{\partial r} + B_z \frac{\partial B_r}{\partial z} - \frac{B_\phi^2}{r} \right) = 0$$

- with  $B_{r,\phi} = 0$

$$\frac{\partial}{\partial r} \left( p + \frac{B^2}{2\mu_0} \right) = 0$$

- and therefore *horizontal pressure balance*

$$p_{\text{inside}} + \frac{B^2}{2\mu_0} = p_{\text{outside}}$$

## Radial Force Balance

- force balance in general coordinate system

$$\nabla \left( \rho + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} \left( \vec{B} \cdot \nabla \right) \vec{B} = \vec{F}_{\text{gravity}}$$

- in cylindrical coordinates, radial component

$$\frac{\partial}{\partial r} \left( \rho + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} \left( B_r \frac{\partial B_r}{\partial r} + B_z \frac{\partial B_r}{\partial z} - \frac{B_\phi^2}{r} \right) = 0$$

- with  $B_{r,\phi} = 0$

$$\frac{\partial}{\partial r} \left( \rho + \frac{B^2}{2\mu_0} \right) = 0$$

- and therefore *horizontal pressure balance*

$$\rho_{\text{inside}} + \frac{B^2}{2\mu_0} = \rho_{\text{outside}}$$



## Radial Force Balance

- force balance in general coordinate system

$$\nabla \left( \rho + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} \left( \vec{B} \cdot \nabla \right) \vec{B} = \vec{F}_{\text{gravity}}$$

- in cylindrical coordinates, radial component

$$\frac{\partial}{\partial r} \left( \rho + \frac{B^2}{2\mu_0} \right) - \frac{1}{\mu_0} \left( B_r \frac{\partial B_r}{\partial r} + B_z \frac{\partial B_r}{\partial z} - \frac{B_\phi^2}{r} \right) = 0$$

- with  $B_{r,\phi} = 0$

$$\frac{\partial}{\partial r} \left( \rho + \frac{B^2}{2\mu_0} \right) = 0$$

- and therefore *horizontal pressure balance*

$$\rho_{\text{inside}} + \frac{B^2}{2\mu_0} = \rho_{\text{outside}}$$

## Vertical Force Balance

- in the z-direction (along the field lines)

$$\frac{\partial p}{\partial z} = -\rho g$$

- with ideal gas law  $\rho = \frac{\mu p}{kT}$

$$\frac{\partial p}{\partial z} = -\frac{\mu g}{kT} p$$

- pressure as a function of height z

$$p(z) = p(z_0) \exp\left(-\int_{z_0}^z \frac{1}{H(z')} dz'\right)$$

- with the *pressure scale height*

$$H(z) = \frac{kT}{\mu g}$$

## Vertical Force Balance

- in the z-direction (along the field lines)

$$\frac{\partial p}{\partial z} = -\rho g$$

- with ideal gas law  $\rho = \frac{\mu p}{kT}$

$$\frac{\partial p}{\partial z} = -\frac{\mu g}{kT} p$$

- pressure as a function of height z

$$p(z) = p(z_0) \exp\left(-\int_{z_0}^z \frac{1}{H(z')} dz'\right)$$

- with the *pressure scale height*

$$H(z) = \frac{kT}{\mu g}$$

## Vertical Force Balance

- in the z-direction (along the field lines)

$$\frac{\partial p}{\partial z} = -\rho g$$

- with ideal gas law  $\rho = \frac{\mu p}{kT}$

$$\frac{\partial p}{\partial z} = -\frac{\mu g}{kT} p$$


- pressure as a function of height z

$$p(z) = p(z_0) \exp\left(-\int_{z_0}^z \frac{1}{H(z')} dz'\right)$$

- with the *pressure scale height*


$$H(z) = \frac{kT}{\mu g}$$

## 2-D Simulations by Oskar Steiner




# Faculae


## The Sun in White Light



2004/02/12 17:36

## The Sun without Limb Darkening






- 3-D impression when looking at images
- Faculae appear predominantly in plages
- Facular brightenings on disk-center side of granules
- Brightening can extend over about 0.5 arcsec
- Narrow, dark lanes centerward of the facular brightening





Observations by Lites et al. (2004)



Simulations by Keller,  
Schüssler, Vögler, Zakharov,  
ApJL 607, L59 (2004 May 20)

Simulation, 200 G




Simulation, 400 G

The diagram illustrates the structure of solar granules. A central gray shaded region represents a magnetic boundary, which is a downward-sloping plane. Below this boundary, the granule tops are shown as curved lines. A bracket on the left indicates that the region between the magnetic boundary and the granule tops contributes 10-90% of the brightness. Dashed lines represent the granule tops in the absence of the magnetic boundary.


**magnetic  
boundary**

**10 - 90% of  
brightness  
contribution**

**granule "top"**





# Observations of the Solar Cycle



## Sunspot Number

- 11-year (average) cycle period
- as short as 8 years
- as long as 15 years
- amplitude is variable
- stronger cycles are shorter
- Maunder minimum is real
- many things correlate with solar cycle


## Hale's Polarity Law



[www.nso.edu](http://www.nso.edu)

- magnetic Carrington maps on 2 July 1988 and 28 May 1999
- bipolar groups have constant magnetic polarity during one cycle
- magnetic polarity is opposite on opposite hemispheres

# Polar Fields




NASA/NSSTC/Hathaway 2005/10

[science.nasa.gov/ssl/pad/solar/dynamo.htm](http://science.nasa.gov/ssl/pad/solar/dynamo.htm)

- polar fields change polarity in synchrony with bipolar regions
- unipolar fields at the poles
- 22-year magnetic cycle (*Hale Cycle*)



## Spörer's Law



[science.nasa.gov/ssl/pad/solar/sunspots.htm](http://science.nasa.gov/ssl/pad/solar/sunspots.htm)

- latitude dependence with cycle noted by Scheiner and Carrington
- studied in detail by Gustav Spörer
- *butterfly diagram*

# Joy's Law




[science.nasa.gov/ssl/pad/solar/dynamo.htm](http://science.nasa.gov/ssl/pad/solar/dynamo.htm)

[www.ociw.edu/ociw/babcock/howardtalk.pdf](http://www.ociw.edu/ociw/babcock/howardtalk.pdf)


- sunspot groups are tilted with respect to equator
- tilt angle depends on latitude (*Joy's Law*)
- leading spots are closer to equator than following





## Other Cycle Indicators

- many solar parameters depend on solar cycle
- emission in chromospheric lines
- radio emission
- cosmic rays




## Long-Term Records

- potential records longer than sunspot observations
- aurorae
- radioactive isotopes due to cosmic rays

# Neutron Flux

NSF ATM 03-39627


The University of New Hampshire/EOS Chicago/LASR Cosmic Physics Instruments in Space




## Cosmic Rays

- cosmic rays: particles originating from outside the Earth's atmosphere, electrically charged, often with high energies, mostly atomic nuclei
- galactic cosmic rays from outside the solar system
- anomalous cosmic rays coming from interstellar space at edge of heliopause
- *Solar Energetic Particles* from solar flares and coronal mass ejections
- galactic cosmic rays produce neutrons in the Earth's atmosphere
- solar cosmic rays rarely have high enough energy to produce neutrons
- solar and Earth's magnetic field deflect cosmic rays
- ⇒ anti-correlation between cosmic ray flux and sunspot cycle

# Sun-Magnetosphere Interaction




## Carbon-14




- mostly produced between 9 and 15 km
- fast oxidation to carbon dioxide
- radioactive  $\text{CO}^2$  part of carbon cycle
- half-life of about 5730 years
- photosynthesis in plants absorbs  $^{14}\text{C}$
- atmospheric  $^{14}\text{C}$  content: equilibrium between production by cosmic rays, radioactive decay, exchange with other reservoirs
- $^{14}\text{C}$  'frozen' into dead plants and decays
- knowing initial  $^{14}\text{C}$  concentration is basis of radiocarbon dating
- calibration with dated material such as tree-rings

# Carbon-14 in Modern Times



# Solar Magnetic Field Relation

## Sunspot activity and $\delta^{14}\text{C}$





## Reconstructed Sunspot Numbers

