

# Interaction of H<sub>2</sub>S with H atoms on grain surfaces under molecular cloud conditions

J. C. Santos, H. Linnartz, and K.-J. Chuang

Laboratory for Astrophysics, Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands  
e-mail: [santos@strw.leidenuniv.nl](mailto:santos@strw.leidenuniv.nl)

Received 3 July 2023 / Accepted 14 August 2023

## ABSTRACT

**Context.** Hydrogen sulfide (H<sub>2</sub>S) is thought to be efficiently formed on grain surfaces through the successive hydrogenation of sulfur atoms. Its non-detection so far in astronomical observations of icy dust mantles thus indicates that effective destruction pathways must play a significant role in its interstellar abundance. While chemical desorption has been shown to remove H<sub>2</sub>S very efficiently from the solid phase, in line with H<sub>2</sub>S gas-phase detections, possible ice chemistry triggered by the related HS radical have been largely disregarded so far, despite it being an essential intermediate in the H<sub>2</sub>S + H reaction scheme.

**Aims.** We aim to thoroughly investigate the fate of H<sub>2</sub>S upon H-atom impact under molecular cloud conditions, providing a comprehensive analysis combined with detailed quantification of both the chemical desorption and ice chemistry that ensues.

**Methods.** We performed experiments in an ultrahigh vacuum chamber at temperatures between 10 and 16 K in order to investigate the reactions between H<sub>2</sub>S molecules and H atoms on interstellar ice analogs. The changes in the solid phase during H-atom bombardment were monitored in situ by means of reflection absorption infrared spectroscopy (RAIRS), and desorbed species were complementarily measured with a quadrupole mass spectrometer (QMS).

**Results.** We confirmed the formation of H<sub>2</sub>S<sub>2</sub> via reactions involving H<sub>2</sub>S + H and quantified its formation cross section under the employed experimental conditions. Additionally, we directly assessed the chemical desorption of H<sub>2</sub>S by measuring the gas-phase desorption signals with the QMS, providing unambiguous desorption cross sections. Chemical desorption of H<sub>2</sub>S<sub>2</sub> was not observed. The relative decrease of H<sub>2</sub>S ices by chemical desorption changed from ~85% to ~74% between temperatures of 10 and 16 K, while the decrease as the result of H<sub>2</sub>S<sub>2</sub> formation was enhanced from ~15% to ~26%, suggesting an increasingly relevant sulfur chemistry induced by HS radicals at warmer environments. The astronomical implications are further discussed.

**Key words.** astrochemistry – methods: laboratory: solid state – infrared: ISM – ISM: molecules

## 1. Introduction

Interstellar dense clouds are known for harboring a lavish chemical inventory spanning from simple ions and radicals to a large variety of complex organic molecules (COMs). At the temperatures and densities typical of these environments ( $T = 10\text{--}20\text{ K}$  and  $\rho = 10^4\text{--}10^5\text{ cm}^{-3}$ , respectively; van Dishoeck et al. 2013), thermal desorption cannot take place, and most species – except for H<sub>2</sub> and He – should be fully depleted into interstellar icy dust grains (Collings et al. 2004). Yet, observations with radio-astronomical facilities have detected copious amounts of COMs such as methanol (CH<sub>3</sub>OH), acetaldehyde (CH<sub>3</sub>CHO), methyl formate (CH<sub>3</sub>OCHO), among others, in the gas phase toward dense and cold clouds (see, e.g., Öberg et al. 2010; Bacmann et al. 2012; Cernicharo et al. 2012; Jiménez-Serra et al. 2016; Scibelli & Shirley 2020). Especially given that these hydrogen-rich species are most likely formed in the ice mantles that shroud interstellar dust grains, such observations reveal that non-thermal desorption mechanisms must play a significant role in balancing gas- and solid-phase chemical abundances. For smaller species, such as CO, photodesorption induced by UV photons through the (in-)direct desorption induced by electronic transitions (DIET) mechanism is an efficient desorption process that could explain, in part, the observed abundances of gaseous species (Öberg et al. 2007; Muñoz Caro et al. 2010; Fayolle et al. 2011; Chen et al. 2014; Paardekooper et al. 2016; Sie et al. 2022).

However, larger molecules are increasingly susceptible to fragmentation upon UV photon impact, which can then be followed by photochemical desorption (Bertin et al. 2016; Cruz-Diaz et al. 2016). Moreover, recent studies have shown that the photodesorption of CO and CH<sub>3</sub>OH ices induced by IR photons might be astronomically relevant (Santos et al. 2023), shedding light on potential new processes to help explain gas-phase abundances of COMs.

Complementarily, another promising non-thermal desorption mechanism that proceeds without fragmentation is the so-called chemical desorption, or reactive desorption. This phenomenon involves the ejection of products upon formation in an exothermic reaction. It has been consistently shown to improve gas-phase abundances predicted by chemical models (Garrod et al. 2006, 2007; Cazaux et al. 2010; Vasyunin & Herbst 2013; Vidal et al. 2017; Cuppen et al. 2017; Fredon et al. 2021) and has been explored in the laboratory for a range of astronomically relevant species and substrates (Dulieu et al. 2013; Minissale & Dulieu 2014; Minissale et al. 2016; He et al. 2017; Chuang et al. 2018; Oba et al. 2018, 2019; Nguyen et al. 2020, 2021). Yet, efforts to experimentally quantify chemical desorption efficiencies are still limited, and modelers typically assume a universal input value between 0.01 and 0.1 (Garrod et al. 2007; Cuppen et al. 2017).

Among the species whose observed abundances cannot be explained by gas-phase processes alone, hydrogen sulfide (H<sub>2</sub>S)

is perhaps one of the most broadly studied in the recent literature. It has been detected toward various interstellar sources and in the comae of comets (Thaddeus et al. 1972; Minh et al. 1989; van Dishoeck et al. 1995; Hatchell et al. 1998; Vastel et al. 2003; Wakelam et al. 2004; Neufeld et al. 2015; Le Roy et al. 2015; Biver et al. 2015; Calmonte et al. 2016; Phuong et al. 2018; Navarro-Almaida et al. 2020). It was also tentatively identified on the surface of the Galilean satellites Io, Ganymede, and Callisto (Nash & Howell 1989; McCord et al. 1998). However, solid-phase interstellar H<sub>2</sub>S has not been unequivocally detected yet, and only upper limits are available in ices so far (Smith 1991; van der Tak et al. 2003; Jiménez-Escobar & Muñoz Caro 2011).

The main proposed route to form H<sub>2</sub>S is through the successive hydrogenation of sulfur on icy grains (S  $\xrightarrow{+H}$  HS  $\xrightarrow{+H}$  H<sub>2</sub>S). Once formed, H<sub>2</sub>S can undergo an H-induced abstraction reaction to form the radical HS



by quantum tunneling through an effective barrier of  $\sim 1500$  K (Lamberts & Kästner 2017). The HS radical can subsequently be hydrogenated to reform H<sub>2</sub>S. Alternatively, H<sub>2</sub>S can also be energetically processed to form species such as H<sub>2</sub>S<sub>2</sub> and a wide range of S allotropes (Moore et al. 2007; Garozzo et al. 2010; Jiménez-Escobar & Muñoz Caro 2011; Jiménez-Escobar et al. 2014; Chen et al. 2015; Shingledecker et al. 2020; Cazaux et al. 2022; Mifsud et al. 2022).

Laboratory studies have reported the hydrogenation of a thin layer (0.7 monolayers; ML) of H<sub>2</sub>S on top of both porous and non-porous amorphous solid water as well as on top of polycrystalline water ice (Oba et al. 2018, 2019). The experimental data has demonstrated that the excess energy generated by the cycle of H-induced abstraction and H<sub>2</sub>S reformation results in chemical desorption with high effectiveness. Kinetic Monte Carlo simulations of such experiments suggest the chemical desorption efficiency to be of  $\sim 3\%$  per hydrogenation event (Furuya et al. 2022). Contrary to energetically processed ices, however, new species formed by the HS radicals were not reported – possibly due to the relatively low abundance of H<sub>2</sub>S species in their experiments. In this work, we aim to further constrain the chemical desorption efficiency of H<sub>2</sub>S by incorporating the chemistry involving HS radicals resulting from the (de-)hydrogenation of hydrogen sulfide, in particular to form disulfane (H<sub>2</sub>S<sub>2</sub>). Moreover, for the first time, we present a comprehensive experimental analysis of the H<sub>2</sub>S chemical desorption phenomenon supported by a strong gas-solid correlation using infrared spectroscopy and mass spectrometry techniques concomitantly.

The experimental setup and techniques employed are described in Sect. 2. The results are shown and discussed in Sect. 3, where we provide effective cross sections for the chemical desorption of H<sub>2</sub>S and H<sub>2</sub>S<sub>2</sub> formation. In Sect. 4, the astrochemical implications of this work are considered, and our main findings are summarized in Sect. 5.

## 2. Experimental methods

We performed our experiments using the ultrahigh vacuum setup SURFRESIDE<sup>3</sup>, which has been described in detail elsewhere (Ioppolo et al. 2013; Qasim et al. 2020). Thus, we only provide the relevant information in this work. The main chamber operates at a base pressure of  $\sim 5 \times 10^{-10}$  mbar. In its center is a gold-plated copper substrate mounted on the tip of a closed cycle He cryostat. The temperature of the substrate can vary

between 8 and 450 K through resistive heating and is monitored by two silicon diode sensors with a relative accuracy of 0.5 K. Ices of H<sub>2</sub>S (Linde, purity 99.5%) are deposited either prior to or simultaneously with H atoms generated by a hydrogen atom beam source (HABS; Tschersich 2000) during what is referred to in this work as pre- and codeposition experiments, respectively. The hydrogen atoms are cooled to room temperature through collision with a nose-shaped quartz pipe before reaching the substrate. As described in detail by Ioppolo et al. (2013), the determination of the absolute H-atom flux is done by placing a quadrupole mass spectrometer (QMS) at the exact position of the substrate and monitoring its signal in a series of systematic experiments with varying filament temperatures and inlet gas flow. Such a measurement is not a trivial procedure, but it serves as a reference guide for regular calibrations of the relative H flux at different operation conditions through the HO<sub>2</sub> peak intensity formed in the barrierless reaction H + O<sub>2</sub> → HO<sub>2</sub>. To infer the temperature-dependent kinetics of the processes explored in this work, we performed predeposition experiments at a range of temperatures of relevance to interstellar molecular clouds (10, 12, 14, and 16 K). Due to its low sticking coefficient at the studied temperatures, the presence of H<sub>2</sub> molecules on the ice (either incoming from the atom source or formed through H recombination) was not expected to significantly affect the outcome of our experiments (Watanabe & Kouchi 2002; Ioppolo et al. 2010).

Ice growth through vapor deposition is monitored by Fourier-transform reflection-absorption infrared spectroscopy (FT-RAIRS). The IR spectra are acquired in the range of 700–4000 cm<sup>-1</sup>, with a resolution of 1 cm<sup>-1</sup>. Concurrently, species in the gas phase are ionized upon electron impact with 70 eV and recorded by a QMS. Once the depositions are finished, temperature-programmed desorption (TPD) experiments are performed by heating the sample at a ramping rate of 5 K min<sup>-1</sup> while concomitantly monitoring the solid and gas phases with the RAIRS and QMS techniques, respectively.

The column densities ( $N_X$ ) of the species in the ice are derived by converting the IR integrated absorbance ( $\int Abs(\nu)d\nu$ ) to absolute abundance using a modified Beer-Lambert law:

$$N_X = \ln 10 \frac{\int Abs(\nu)d\nu}{A'(X)}, \quad (2)$$

where  $A'(X)$  is the apparent absorption band strength of a given species. For H<sub>2</sub>S, band strength values measured by infrared transmission spectroscopy are available in the literature. However, signals obtained in reflection mode are systematically higher than transmission counterparts due to substrate dipole couplings and a typically longer IR pathway in the ice. Thus, to ensure high accuracy in the derivation of the H<sub>2</sub>S ice column density, we performed calibration experiments using the laser interference technique that yield a band strength value of  $A'(\text{H}_2\text{S})_{\sim 2553 \text{ cm}^{-1}} \sim (4.7 \pm 0.1) \times 10^{-17} \text{ cm molecule}^{-1}$  for our specific experimental settings (see Appendix A).

Since direct determination of the H<sub>2</sub>S<sub>2</sub> band strength is challenging, we estimate  $A'(\text{H}_2\text{S}_2)$  in a similar way to what has been described by Cazaux et al. (2022). The column density ratio ( $N_{\text{H}_2\text{S}_2}/N_{\text{H}_2\text{S}}$ ) can be derived from the QMS data by the expression (Martín-Doménech et al. 2015)

$$\frac{N_{\text{H}_2\text{S}_2}}{N_{\text{H}_2\text{S}}} = \frac{A(66)}{A(34)} \cdot \frac{\sigma^+(\text{H}_2\text{S})}{\sigma^+(\text{H}_2\text{S}_2)} \cdot \frac{I_F([\text{H}_2\text{S}]^+)}{I_F([\text{H}_2\text{S}_2]^+)}. \frac{F_F(34)}{F_F(66)} \cdot \frac{S(34)}{S(66)}, \quad (3)$$

where  $A(m/z)$  is the integrated area of a given mass fragment;  $\sigma^+(X)$  is the molecule's electronic ionization cross section;  $I_F(z)$

**Table 1.** List of parameters used in the estimation of  $A'(\text{H}_2\text{S}_2)$ .

Species	$\alpha$ ( $\text{\AA}^3$ ) <sup>(a)</sup>	$F_F(m/z)$ <sup>(b)</sup>	$S(m/z)$ <sup>(b)</sup>
H <sub>2</sub> S	3.776	0.52	0.28
H <sub>2</sub> S <sub>2</sub>	6.828	0.31	0.08

**Notes.** <sup>(a)</sup>CCCBDB. <sup>(b)</sup>Values are given for the molecular ions.

is the ionization fraction of charge  $z$  (in this work, corresponding to unity);  $F_F(m/z)$  is the fragmentation fraction; and  $S(m/z)$  is the sensitivity of the QMS at a specific mass. As there are no values for  $\sigma^+(\text{H}_2\text{S}_2)$  reported in the literature, we estimated its value based on the molecule's polarizability volume ( $\alpha(X)$ ) by the empirical correlation (Hudson et al. 2006; Bull et al. 2012):

$$\sigma_{\text{max}}^+(X) = c \cdot \alpha(X), \quad (4)$$

where  $X$  denotes a given species and  $c$  is a correlation constant of  $1.48 \text{ \AA}^{-1}$ . The maximum ionization cross section ( $\sigma_{\text{max}}^+$ ) of organic species typically occurs around 90 eV and varies only slightly ( $< 5\%$ ) in intensity from ionizations with 70 eV (Hudson et al. 2003; Bull & Harland 2008). Thus, we utilized this method to derive both  $\sigma^+(\text{H}_2\text{S}_2)$  and  $\sigma^+(\text{H}_2\text{S})$  from  $\alpha(\text{H}_2\text{S}_2)$  and  $\alpha(\text{H}_2\text{S})$  as calculated by group additivity<sup>1</sup>. The  $F_F(m/z)$  of the relevant mass fragments are inferred from the QMS data acquired during the TPD experiments after codeposition of H<sub>2</sub>S and H, and we obtained the sensitivity from previous calibrations performed with the same setup (Chuang 2018). The employed values are summarized in Table 1.

By combining  $(N_{\text{H}_2\text{S}_2})/(N_{\text{H}_2\text{S}})$  from Eq. (3) and  $N_{\text{H}_2\text{S}}$  from Eq. (2), one can obtain  $N_{\text{H}_2\text{S}_2}$ , which in turn can be used to estimate  $A'(\text{H}_2\text{S}_2)$  from the integrated absorbance area of the IR spectra:

$$A'(\text{H}_2\text{S}_2) = \frac{\int Abs(\nu) d\nu}{N_{\text{H}_2\text{S}_2}}. \quad (5)$$

The average between two independent experiments yields an estimated  $A'(\text{H}_2\text{S}_2)_{\sim 2490 \text{ cm}^{-1}} \sim (9.9 \pm 0.2) \times 10^{-17} \text{ cm molecule}^{-1}$ .

The details of the experiments performed in this work are summarized in Table 2. The relative errors of both H<sub>2</sub>S and H fluxes are estimated to be  $\sim 5\%$ .

### 3. Results and discussion


#### 3.1. H<sub>2</sub>S + H ice chemistry

The left panel of Fig. 1 shows the spectra obtained after deposition of pure H<sub>2</sub>S and codeposition of H<sub>2</sub>S + H at 10 K in the frequency region characteristic of SH stretching modes. A strong IR feature is observed at  $\sim 2553 \text{ cm}^{-1}$ , corresponding to the  $\nu_1$  (symmetric) and  $\nu_3$  (anti-symmetric) SH stretching modes of H<sub>2</sub>S. In comparison, when H atoms are also present, a new feature peaking at  $\sim 2490 \text{ cm}^{-1}$  appears on the red wing of the  $\nu_{1,3}$  mode of H<sub>2</sub>S – consistent with the SH stretching band ( $\nu_1$ , sym.; and  $\nu_5$ , anti-sym.) of H<sub>2</sub>S<sub>2</sub> (Isoniemi et al. 1999). During the TPD experiment performed after codepositing H<sub>2</sub>S + H, the main bands at  $\sim 2553 \text{ cm}^{-1}$  and  $\sim 2490 \text{ cm}^{-1}$  fully disappear in

<sup>1</sup> Values have been taken from the NIST Computational Chemistry Comparison and Benchmark Database (CCCBDB), NIST Standard Reference Database Number 101, <https://cccbdb.nist.gov/>

**Table 2.** Overview of the experiments performed in this work.

Experiment	$T_{\text{sample}}$ (K)	H <sub>2</sub> S flux ( $\text{cm}^{-2} \text{ s}^{-1}$ )	H flux ( $\text{cm}^{-2} \text{ s}^{-1}$ )	Time (min)
Codeposition experiments				
H <sub>2</sub> S	10	$\sim 1 \times 10^{13}$		60
H <sub>2</sub> S + H	10	$\sim 1 \times 10^{13}$	$\sim 8 \times 10^{12}$	60
Predeposition experiments				
H <sub>2</sub> S $\rightarrow$ +H	10	$\sim 1 \times 10^{13}$	$\sim 8 \times 10^{12}$	60 + 120
H <sub>2</sub> S $\rightarrow$ +H	12	$\sim 1 \times 10^{13}$	$\sim 8 \times 10^{12}$	60 + 120
H <sub>2</sub> S $\rightarrow$ +H	14	$\sim 1 \times 10^{13}$	$\sim 8 \times 10^{12}$	60 + 120
H <sub>2</sub> S $\rightarrow$ +H	16	$\sim 1 \times 10^{13}$	$\sim 8 \times 10^{12}$	60 + 120




**Fig. 1.** Infrared spectra of the codeposition experiments. Panel a: comparison between the final infrared spectra after deposition of a pure H<sub>2</sub>S ice (black) superimposed by the final spectrum after codeposition of H<sub>2</sub>S and H atoms (red) with analogous experimental conditions. Panel b: infrared spectra acquired during the warming up of the H<sub>2</sub>S ice codeposited with H atoms, offset for clarity. In both panels, the assignments of the H<sub>2</sub>S and H<sub>2</sub>S<sub>2</sub> vibrational bands are shown with dashed lines.


the temperature ranges of 10–100 K and 100–140 K, respectively (Fig. 1, right panel), which coincides with previously measured desorption temperatures of H<sub>2</sub>S and H<sub>2</sub>S<sub>2</sub> (Jiménez-Escobar & Muñoz Caro 2011; Chen et al. 2015; Cazaux et al. 2022).

The assignments of H<sub>2</sub>S and H<sub>2</sub>S<sub>2</sub> are substantiated by their respective mass fragments induced by electron impact during the TPD experiments in Figs. 2 and 3, respectively. As shown in Fig. 2a, a desorption peak of fragments  $m/z = 32$  and  $34$  can be observed at  $\sim 85 \text{ K}$  in both H<sub>2</sub>S and H<sub>2</sub>S + H cases, amounting to relative intensities consistent with the standard for H<sub>2</sub>S as provided by the NIST database<sup>2</sup>. This desorption temperature matches the disappearance of the  $\sim 2553 \text{ cm}^{-1}$  bands in the IR spectra. In Fig. 3, the desorption peak of the mass fragments associated with H<sub>2</sub>S<sub>2</sub> is detected solely in the H<sub>2</sub>S + H experiment at 126 K – coinciding with the disappearance of the feature at  $\sim 2490 \text{ cm}^{-1}$  in the IR spectra. Thus, the assignment of the new peak as H<sub>2</sub>S<sub>2</sub> is confirmed by both RAIRS and QMS techniques combined with TPD experiments. Given the lack of laboratory data on its mass fragmentation pattern, we provide for the first

<sup>2</sup> <https://webbook.nist.gov/chemistry/>





**Fig. 2.** TPD-QMS results for the detection of  $\text{H}_2\text{S}$ . Panel a: TPD-QMS spectra of  $m/z = 32$  (blue) and  $m/z = 34$  (red) after deposition of a pure  $\text{H}_2\text{S}$  ice and codeposition of  $\text{H}_2\text{S} + \text{H}$  with analogous experimental conditions. Spectra are offset for clarity and shown in the temperature range relevant to  $\text{H}_2\text{S}$  thermal desorption. Panel b: Comparison between the relative intensities of  $m/z = 32$  and  $34$  desorbing at  $85\text{ K}$  in both  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S} + \text{H}$  experiments together with the standard fragmentation pattern of  $\text{H}_2\text{S}$  from NIST.



**Fig. 3.** TPD-QMS results for the detections of  $\text{H}_2\text{S}_2$ . Panel a: TPD-QMS spectra of  $m/z = 32$  (blue),  $34$  (red),  $64$  (green),  $65$  (purple), and  $66$  (yellow) after deposition of a pure  $\text{H}_2\text{S}$  ice and codeposition of  $\text{H}_2\text{S} + \text{H}$  with analogous experimental conditions. Spectra are offset for clarity and shown in the temperature range relevant to  $\text{H}_2\text{S}_2$  thermal desorption. Panel b: mass fragmentation pattern of  $\text{H}_2\text{S}_2$  generated by  $70\text{ eV}$  electron ionization as measured in this work.

time – to the best of our knowledge – the relative intensities of  $m/z = 32, 34, 64, 65,$  and  $66$  as generated by  $70\text{ eV}$  electron ionization of  $\text{H}_2\text{S}_2$  and corrected for the sensitivity of the QMS in the right panel of Fig. 3. The contribution from the  $^{34}\text{S}$  isotope (natural abundance of  $4.29\%$ ) is included in the fragmentation pattern.

When  $\text{H}_2\text{S}$  is deposited simultaneously with  $\text{H}$  atoms,  $\text{HS}$  radicals formed by the hydrogen abstraction of  $\text{H}_2\text{S}$  (Reaction (1)) can thus further associate either with  $\text{H}$  atoms, reforming  $\text{H}_2\text{S}$ , or with  $\text{HS}$  radicals, forming  $\text{H}_2\text{S}_2$ :



**Fig. 4.** Infrared spectrum after deposition of a pure  $\text{H}_2\text{S}$  ice (black) and the difference spectra after exposure to  $\text{H}$  atoms for  $20\text{ min}$  (red),  $40\text{ min}$  (blue), and  $60\text{ min}$  (green). Superimposed onto the difference spectra are the corresponding Gaussian fittings of the  $\text{H}_2\text{S}$  band (purple),  $\text{H}_2\text{S}_2$  band (yellow), and the resulting convoluted feature (brown). The spectra have been offset for clarity.


Reaction (6a) proceeds barrierlessly and can result in chemical desorption due to its high exothermicity ( $\sim 45\,000\text{ K}$ , based on the gas-phase enthalpies of formation of reactants and products). Reaction (6b) is also barrierless and has been proposed in previous studies on the energetic processing of  $\text{H}_2\text{S}$ -containing ices (Jiménez-Escobar & Muñoz Caro 2011; Jiménez-Escobar et al. 2014; Chen et al. 2015; Cazaux et al. 2022; Mifsud et al. 2022).

### 3.2. H-atom bombardment on $\text{H}_2\text{S}$ ice

The left panels in Figs. 1 and 2 show that the amount of  $\text{H}_2\text{S}$  ice after the codeposition experiment with  $\text{H}$  atoms is smaller than that of the pure ice deposition with the same experimental conditions, thus signaling that the interaction of  $\text{H}_2\text{S}$  with hydrogen leads to a net loss of material as a result of both Reactions (6a) and (6b). While the efficiency of the former reaction has been explored in detail (Oba et al. 2018, 2019; Furuya et al. 2022), the contribution from  $\text{H}_2\text{S}_2$  formation to depleting  $\text{H}_2\text{S}$  from the solid phase has not been considered so far. In this work, we explore the effectiveness of both reactions thoroughly and assess their respective relevance to the destruction of the  $\text{H}_2\text{S}$  ice.

To quantify the efficiencies of Reactions (6a) and (6b), we monitored the abundance of  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_2$  as a function of  $\text{H}$ -atom fluence during predeposition experiments in which a deposited  $\text{H}_2\text{S}$  ice was subsequently bombarded by a constant  $\text{H}$ -atom flux. The difference spectra after the  $\text{H}$ -atom bombardment for  $20, 40,$  and  $60\text{ min}$  at  $10\text{ K}$  together with the pure  $\text{H}_2\text{S}$  sample prior to hydrogenation are shown in Fig. 4. Both  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_2$  features can be resolved in the difference spectra by deconvolution using Gaussian profiles, as shown by the superimposing lines. The interaction with  $\text{H}$  atoms leads to a loss of  $\text{H}_2\text{S}$ , as evinced by the decrease in its  $\text{SH}$  stretching band at  $\sim 2553\text{ cm}^{-1}$  (purple dashed line). Concomitantly, a feature due






**Fig. 5.** Monitoring of H<sub>2</sub>S in the solid and gas phases during H-atom bombardment. Upper panel: variation in H<sub>2</sub>S column density measured from the  $\sim 2553\text{ cm}^{-1}$  band in the IR spectra as a function of time. Lower panel: scan of the  $m/z = 34$  ([H<sub>2</sub>S]<sup>+</sup>) as measured by the QMS as a function of time. The shadowed areas denote the periods during which the H-atom flux was stopped.

to the SH stretching modes of H<sub>2</sub>S<sub>2</sub> appears on the red wing of the H<sub>2</sub>S band and becomes increasingly evident at longer H-atom exposure times (yellow dashed line). The results of the predeposition experiments are therefore consistent with the codeposition counterparts, and they indicate a non-negligible contribution to the H<sub>2</sub>S depletion from Reaction (6b). In contrast, neither Oba et al. (2018) nor Oba et al. (2019) have detected any other sulfur-bearing species apart from hydrogen sulfide during similar H<sub>2</sub>S → +H predepositions at 10–30 K followed by TPD experiments. Such a discrepancy might be due to the limited abundance of H<sub>2</sub>S in the aforementioned works (0.7 ML) compared to the present experiments ( $\sim 40$  ML), as they might not yield product amounts above the instrumental detection limit.

To directly probe the chemical desorption of H<sub>2</sub>S as a result of reactions with H atoms, we monitored the gas-phase signals via the relevant mass fragments ( $m/z = 34$ , [H<sub>2</sub>S]<sup>+</sup>;  $m/z = 33$ , [HS]<sup>+</sup>) with a QMS during the H-exposure experiments. In Fig. 5, data acquired by both the RAIRS and QMS techniques while intermittently bombarding the predeposited H<sub>2</sub>S ice with H atoms (i.e., in three intervals of 20 min, ending with 60 min) are presented in the upper and lower panels, respectively. In the first 20 min of bombardment, we observed a steep decrease in the H<sub>2</sub>S IR absorbance area coinciding with an abrupt increase in the  $m/z = 34$  readout by the QMS. Once bombardment was stopped, the area of the H<sub>2</sub>S band remained fairly constant, and the QMS signal dropped to the base value. Such results provide unambiguous evidence of the effective chemical desorption of H<sub>2</sub>S upon H-atom exposure. Following the first bombardment, a similar behavior was observed through both RAIRS and QMS techniques for the rest of the exposure periods, albeit to a diminishing extent of H<sub>2</sub>S loss due to saturation of the




**Fig. 6.** Derived contributions from H<sub>2</sub>S<sub>2</sub> formation and H<sub>2</sub>S chemical desorption to the measured loss in  $N(\text{H}_2\text{S})$  after 120 min of H-atom exposure at 10, 12, 14, and 16 K.

ice layer within the penetration depth of the hydrogen atoms – typically of a few monolayers (see, e.g., Watanabe & Kouchi 2008; Fuchs et al. 2009). No increase in signal was detected for  $m/z = 66$  ([H<sub>2</sub>S<sub>2</sub>]<sup>+</sup>), indicating that relative to H<sub>2</sub>S, disulfane does not undergo chemical desorption effectively upon formation. This is quite expected, as H<sub>2</sub>S<sub>2</sub> contains more degrees of freedom and, as inferred from its higher desorption temperature, a higher binding energy than H<sub>2</sub>S. Consequently, H<sub>2</sub>S<sub>2</sub> does not contribute significantly to the measurement of  $m/z = 34$  during H-atom exposure, which can therefore be solely attributed to H<sub>2</sub>S.

We measured the intensity of the  $m/z = 33$  signal relative to  $m/z = 34$  to be  $\sim 0.55$  throughout the H-atom exposure, whereas the expected fragmentation pattern of H<sub>2</sub>S corresponds to  $33/34 \sim 0.42$ . The excess of [HS]<sup>+</sup> fragments detected during the bombardment is consistent with the transfer of HS radicals to the gas phase through chemical desorption as a result of Reaction (1). This fraction, however, is significantly smaller than the detected gaseous H<sub>2</sub>S and could therefore be neglected. Indeed, due to the high exothermicity of Reaction (6a) and the fact that its excess energy is concentrated in a single product, we expected H<sub>2</sub>S to be the most susceptible species to chemical desorption during the hydrogenation sequence – as was also suggested by Oba et al. (2018).

In addition to experiments at 10 K, we performed predeposition experiments with analogous conditions at 12 K, 14 K, and 16 K in order to investigate the effects of different temperatures on H<sub>2</sub>S<sub>2</sub> formation and H<sub>2</sub>S chemical desorption. The percentage of H<sub>2</sub>S lost either to chemical desorption or H<sub>2</sub>S<sub>2</sub> formation by the end of the predeposition experiments can be derived by comparing the final  $\Delta N$  of both species, assuming that other potential processes have a minor contribution in decreasing the H<sub>2</sub>S band. The derived efficiencies are temperature dependent, as shown in Fig. 6; the overall H<sub>2</sub>S loss due to chemical desorption varies from  $\sim 85\%$  to  $\sim 74\%$  when the ice temperature



**Fig. 7.** Column densities of  $\text{H}_2\text{S}$  and  $\text{H}_2\text{S}_2$  in the ice during H-atom bombardment. Upper panel: variation in  $\text{H}_2\text{S}_2$  column density during H-atom exposure of an  $\text{H}_2\text{S}$  ice at 10 K. Lower panel: variation in  $\text{H}_2\text{S}$  column density as a function of H-atom fluence during bombardment of an  $\text{H}_2\text{S}$  ice at 10 K. The two-term exponential fitting to the points is shown in red, while the fast and slow components of the fitting are plotted in blue and green, respectively.

increases from 10 K to 16 K. Accordingly, the percentage loss due to the formation of  $\text{H}_2\text{S}_2$  varies from  $\sim 15\%$  to  $\sim 26\%$ . We note that these values are respective to the relative  $\text{H}_2\text{S}$  loss at each specific temperature and are not the absolute amount of formed  $\text{H}_2\text{S}_2$  or chemically desorbed  $\text{H}_2\text{S}$  in each experiment. At higher temperatures, the fraction of  $\text{H}_2\text{S}$  consumed to form  $\text{H}_2\text{S}_2$  increases relatively to the loss due to chemical desorption, suggesting that the former process becomes increasingly relevant in warmer environments. This observation is possibly related to a significant increase in diffusion rates of HS radicals enhancing the overall  $\text{H}_2\text{S}_2$  formation at the expense of chemical desorption by  $\text{H}_2\text{S}$  reformation. In summary, by taking into account this chemical loss channel, it is possible to further constrain the fate of  $\text{H}_2\text{S}$  molecules upon H-atom bombardment, thus expanding the results from previous works in which  $\text{H}_2\text{S}_2$  formation was not observed.

### 3.3. Kinetic analysis

Information on the kinetics of  $\text{H}_2\text{S}_2$  formation and  $\text{H}_2\text{S}$  consumption can be derived from predeposition experiments. In the upper panel of Fig. 7, the variation in column density ( $\Delta N$ ) of  $\text{H}_2\text{S}_2$  as a function of H-atom fluence measured from the IR spectra at 10 K is shown. The curve is fitted by a single exponential function:

$$\Delta[X]_t = [\text{H}_2\text{S}]_0 \cdot a(1 - \exp(-\sigma \cdot F)), \quad (7)$$

where  $\Delta[X]$  and  $[\text{H}_2\text{S}]_0$  are, respectively, the abundance of species  $X$  at a given time and the initial abundance of  $\text{H}_2\text{S}$ . Here,  $a$  is the saturation value,  $F$  is the incident H-atom fluence, and

$\sigma$  is the effective formation cross section of  $\text{H}_2\text{S}_2$ . From this fitting, we derived  $\sigma \sim (9.8 \pm 0.9) \times 10^{-17} \text{ cm}^2$  for  $\text{H}_2\text{S}_2$  formation at 10 K. We note, however, that the rate law of  $\text{H}_2\text{S}_2$  formation is far from trivial, as both Reactions (1) and (6b) contribute to the effective cross section, with the latter requiring two HS radicals to occur. Therefore, it cannot be simplified by the pseudo first-order approximation. Moreover, the accurate amount of H atoms available on the surface of the ice is very difficult to quantify, as a fraction will recombine to form  $\text{H}_2$  – hence the use of the “effective” term. The  $\sigma$  value derived here is thus not suited to be directly employed in chemical models as a rate constant, but it is very useful for comparison purposes with other effective cross sections derived with similar conditions.

In the lower panel of Fig. 7, the effective variation in the column density of  $\text{H}_2\text{S}$  as a function of H-atom fluence measured from the infrared spectra is shown. In this case, the plot is better fitted by a two-term exponential function:


$$\Delta[\text{H}_2\text{S}]_t = [\text{H}_2\text{S}]_0(a_1(1 - \exp(-\sigma_1 \cdot F)) + a_2(1 - \exp(-\sigma_2 \cdot F))), \quad (8)$$

where  $a_n$  is the saturation value and  $\sigma_n$  is the effective destruction cross section. The interpretation of such a fitting is not straightforward, as it incorporates the contribution from all the processes leading to a decrease in  $N(\text{H}_2\text{S})$ . Nonetheless, the double exponential fitting suggests that the processes dominating the observed decrease in  $N(\text{H}_2\text{S})$  can be separated into two different timescales with  $\sigma_1 \sim 10^{-16} \text{ cm}^2$  and  $\sigma_2 \sim 10^{-17} \text{ cm}^2$ .

The fast process with  $\sigma_1 \sim 10^{-16} \text{ cm}^2$  is likely due to startup effects, such as collision-induced desorption of the weakly bound topmost molecules (Chuang et al. 2018). Accordingly, the effective destruction cross section of  $\text{H}_2\text{S}$  can be approximated as the second exponential term with  $\sigma_2 \sim 10^{-17} \text{ cm}^2$ . Control experiments with neutral helium bombardment of  $\text{H}_2\text{S}$  ices show that material loss due to collisional impact should account for  $\lesssim 10\%$  of the total  $\text{H}_2\text{S}$  desorption from the QMS. In comparison, the saturation point of the fast exponential curve (blue line in the lower panel of Fig. 7) corresponds to  $\sim 0.3$  of the total loss of  $\text{H}_2\text{S}$  and should thus be regarded as an upper limit to the real value.

Given that the interaction of  $\text{H}_2\text{S}$  with H atoms mostly results in chemical desorption via Reaction (6a) and  $\text{H}_2\text{S}_2$  formation via Reaction (6b), it is possible to isolate the  $\text{H}_2\text{S}$  chemical desorption curve by subtracting the minimum amount of  $\text{H}_2\text{S}$  consumed to form  $\text{H}_2\text{S}_2$  (i.e., twice the column density of  $\text{H}_2\text{S}_2$ ). The resulting isolated  $\text{H}_2\text{S}$  chemical desorption curve is shown in the upper panel of Fig. 8 and yields an effective cross section of  $\sigma \sim (1.7 \pm 0.2) \times 10^{-17} \text{ cm}^2$ . We emphasize, however, that this value was derived using a series of assumptions and is therefore only a rough estimation.

In addition to the IR approach, it is possible to directly probe the chemical desorption of hydrogen sulfide by utilizing mass spectrometry data acquired during hydrogen exposure. The lower panel of Fig. 8 shows the integrated signal for the  $m/z = 34$  ( $[\text{H}_2\text{S}]^+$ ) fragment as a function of H-atom fluence (i.e., the area of the plot in the lower panel of Fig. 5). Similar to  $\text{H}_2\text{S}_2$ , this curve can be fitted by an exponential function as described in Eq. (7), yielding  $\sigma \sim (3.7 \pm 0.3) \times 10^{-17} \text{ cm}^2$ , which is quite compatible with the IR approach. Assuming similar chemical desorption efficiencies for both  $^{32}\text{S}$  and  $^{34}\text{S}$  isotopes of  $\text{H}_2\text{S}$ , the contribution from  $^{34}\text{S}$  to  $m/z = 34$  does not affect the exponential factor in the fitting and can therefore be neglected. We find it important to note that the cross section from the QMS data is likely more accurate than the IR counterpart, as the former is a



**Fig. 8.** Chemical desorption curves for  $\text{H}_2\text{S}$  as a function of H-atom fluence. Upper panel: estimated contribution from chemical desorption to the decrease in  $N(\text{H}_2\text{S})$  as a function of fluence. The simple exponential fitting to the points is shown in red, and the linear fitting to the first 55 min of bombardment is shown in blue (dashed line). Lower panel: integrated intensity of the  $m/z=34$  signal measured by the QMS as a function of H-atom fluence during the same experiment. The red line shows the exponential fitting to the points.

direct fitting of the measurements, whereas the latter involves a number of presumptions. Both values are similar to the chemical desorption cross sections of  $(2.1 \pm 0.2) \times 10^{-17} \text{ cm}^2$  derived by Oba et al. (2019) from the exposure of  $\text{H}_2\text{S}$  ice to H atoms at 10 K and reinforce the relevance of  $\text{H}_2\text{S}$  chemical desorption to interstellar gas-grain chemistry. Small discrepancies between the two studies are expected due to the different experimental conditions, such as ice thicknesses, growth surfaces, and H-atom fluxes.

We performed similar experiments at 12, 14, and 16 K, and the derived effective cross sections are summarized in Table 3. The estimated  $\sigma(\text{H}_2\text{S}_2)$  values suggest that the effectiveness of  $\text{H}_2\text{S}_2$  formation remains fairly consistent (within the uncertainty range) for temperatures between 10 K and 14 K. At 16 K, the cross section is slightly reduced. This behavior is likely the outcome of competing elementary processes involved in synthesizing  $\text{H}_2\text{S}_2$  on ice. While diffusion can be facilitated at higher temperatures, thus enhancing encounters between two HS radicals and favoring Reaction (6b), the sticking coefficient of H atoms on ices diminishes, thus hindering the formation of reactants in the first place. Moreover, faster diffusion rates also imply that H atoms might not have enough available time in the vicinity of an  $\text{H}_2\text{S}$  molecule to overcome the  $\sim 1500$  K barrier in Reaction (1). Similar findings were described in other H-atom addition experiments (e.g., in the hydrogenation of  $\text{O}_2$ ; Ioppolo et al. 2008, 2010; Cuppen et al. 2010).

We obtained the effective cross sections of  $\text{H}_2\text{S}$  chemical desorption from the QMS data, and they show a slight decreasing trend between the temperatures of 10, 12, 14, and 16 K. A similar behavior was also observed by Oba et al. (2019) with

**Table 3.** Effective cross sections of  $\text{H}_2\text{S}_2$  formation ( $\sigma(\text{H}_2\text{S}_2)$ ) and  $\text{H}_2\text{S}$  chemical desorption ( $\sigma_{CD}(\text{H}_2\text{S})$ ) derived from the predeposition experiments performed at 10, 12, 14, and 16 K.

Temperature (K)	$\sigma(\text{H}_2\text{S}_2)$ ( $\times 10^{-17} \text{ cm}^2$ )	$\sigma_{CD}(\text{H}_2\text{S})$ ( $\times 10^{-17} \text{ cm}^2$ )
10	$9.8 \pm 0.9$	$3.7 \pm 0.3$
12	$7.8 \pm 0.9$	$2.8 \pm 0.1$
14	$8.3 \pm 0.7$	$2.7 \pm 0.2$
16	$5.2 \pm 0.6$	$2.6 \pm 0.2$

measurements at 10, 20, and 30 K, which they attributed to the combination of the H atom availability at  $T \geq 20$  K and the true efficiency of  $\text{H}_2\text{S}$  chemical desorption at higher temperatures. The slightly lower effective cross sections, they argue, would in reality indicate an increase of the true value at warmer environments, balancing out the considerably diminishing sticking coefficient of H. In the present work, we probe a much smaller temperature range where the availability of H atoms on the surface is not expected to drop as significantly. Nonetheless, some effect of the smaller sticking coefficient of hydrogen at higher temperatures could in principle influence the measured effective cross sections – albeit to a smaller extent than in Oba et al. (2019). Although speculating the effect of the ice temperature on the real  $\sigma_{CD}(\text{H}_2\text{S})$  is challenging, it seems as though a measurable change occurs only from 10 K to 12 K within the range explored in this work.

#### 4. Astrophysical implications

Hydrogen sulfide is thought to be efficiently formed on the surface of interstellar dust through the hydrogenation of S atoms (see, e.g., Tielens & Hagen 1982; Laas & Caselli 2019). It is also the major sulfur-bearing species found in the comae of comets (Calmonte et al. 2016 and references therein), which in turn are thought to harbor the content of prestellar ices. The (so far) non-detection of solid-phase  $\text{H}_2\text{S}$  in interstellar clouds thus poses a question regarding the fate of  $\text{H}_2\text{S}$  in interstellar icy mantles. One likely explanation for its absence in observations is that solid-phase  $\text{H}_2\text{S}$  is effectively destroyed by, for instance, energetic processing, which is known to result in solid-phase sulfur chemistry (e.g., Moore et al. 2007; Garozzo et al. 2010; Jiménez-Escobar & Muñoz Caro 2011; Jiménez-Escobar et al. 2014; Chen et al. 2015; Shingledecker et al. 2020; Cazaux et al. 2022; Mifsud et al. 2022). In fact, the photochemistry of  $\text{H}_2\text{S}$  induced by UV photons has been suggested as a potential sulfur sink, as it has been shown to produce allotropic forms of S ( $\text{S}_n$ ) that are largely refractory (especially for  $n > 4$ ). In addition to energetic processing, non-energetic routes to remove  $\text{H}_2\text{S}$  from the solid phase are also essential, as they are the dominant processes taking place within dense clouds. Indeed, recent observations with the *James Webb* Space Telescope aimed at highly shielded regions within interstellar clouds (with  $A_V > 50$ ) have still been unable to detect  $\text{H}_2\text{S}$  ices, providing only upper limits of 0.6% with respect to  $\text{H}_2\text{O}$  (McClure et al. 2023). Especially in such environments, chemical desorption due to hydrogenation seems to be a particularly prominent mechanism to transfer  $\text{H}_2\text{S}$  to the gas phase (Oba et al. 2018, 2019). In this work, the cross sections derived directly from the chemically desorbed  $\text{H}_2\text{S}$  as measured by the QMS – and thus not influenced by additional  $\text{H}_2\text{S}$  destruction phenomena such as chemical reactions – is fully in line with this proposition.

Another relevant value that can be derived from predeposition experiments is the efficiency of chemical desorption per incident H atom. The reason for deriving a value per incident atom instead of per reactive event is because the true value of H atoms involved in the reactions under our experimental conditions is unknown, as a fraction of them will recombine into H<sub>2</sub> molecules through diffusion. The efficiency derived per incident atom can therefore be regarded as a lower limit to the value per reaction event. After isolating the variation in the H<sub>2</sub>S column density due to chemical desorption (as described in Sect. 3.3; see also Fig. 8), a linear fit to the points within the first 55 min of bombardment at 10 K (blue dashed line in the lower panel of Fig. 8) yielded an efficiency of  $\sim 0.019 \pm 0.001$ , which is around four times higher than the values reported by Oba et al. (2018, 2019) and consistent with the calculated value per reaction event (i.e.,  $(3 \pm 1.5)\%$ ) in Furuya et al. (2022). Similar to the cross sections, such a discrepancy could be due to the different ice compositions (pure H<sub>2</sub>S versus H<sub>2</sub>S on top of amorphous solid water) and thicknesses ( $\sim 40$  ML versus 0.7 ML). Nonetheless, this estimated efficiency reinforces the key role of H<sub>2</sub>S chemical desorption as a non-thermal mechanism of transferring hydrogen sulfide to the gas phase within dark clouds. Indeed, by combining gas-grain chemical models with millimeter observations, Navarro-Almáida et al. (2020) found that chemical desorption is the main mechanism responsible for gas-phase H<sub>2</sub>S formation.

Complementary to chemical desorption, the interaction of H<sub>2</sub>S with H atoms can also kick-start non-energetic chemistry to form larger sulfur-bearing molecules. The detection of H<sub>2</sub>S<sub>2</sub> under our experimental conditions is one example of how HS radicals produced by Reaction (1) can lead to a higher sulfur-bearing chemical complexity. In fully representative interstellar ices, the probability of two HS radicals meeting is rather low, given the small abundance of H<sub>2</sub>S relative to other ice components, such as H<sub>2</sub>O or CO. However, these radicals can react with more widespread ice species, potentially leading to the formation of sulfur-bearing COMs. This present work therefore serves as a proof of concept that non-energetic sulfur chemistry can be initiated by the formation of HS radicals through Reaction (1) with the simplest example of H<sub>2</sub>S<sub>2</sub>. We find that it is also noteworthy that the contributions from each process to the consumption of H<sub>2</sub>S varies significantly with temperature, with an appreciable increase in sulfur-bearing species formed at 16 K compared to 10 K. This is likely due to the enhanced radical diffusion within warmer ices and signifies that sulfur chemistry could be significantly intensified at regions closer to the edges of dark clouds – where temperatures can approach 20 K.

## 5. Conclusions

In the present work, we experimentally investigated the interaction of H<sub>2</sub>S ices with H atoms under ultrahigh vacuum pressures and astronomically relevant temperatures (10–16 K). Our main findings are summarized below:

- We verified that solid-phase hydrogen sulfide is destroyed and H<sub>2</sub>S<sub>2</sub> is formed as a result of the interaction between H<sub>2</sub>S and H atoms.
- The chemical desorption of H<sub>2</sub>S was directly probed by quantifying the material ejected into the gas phase during H-atom exposure experiments. The calculated effective cross sections for ice temperatures of 10, 12, 14, and 16 K are, respectively,  $(3.7 \pm 0.3) \times 10^{-17} \text{ cm}^2$ ,  $(2.8 \pm 0.1) \times 10^{-17} \text{ cm}^2$ ,  $(2.7 \pm 0.2) \times 10^{-17} \text{ cm}^2$ , and  $(2.6 \pm 0.2) \times 10^{-17} \text{ cm}^2$ .
- From the RAIRS data, we estimate the chemical desorption efficiency per incident H atom at 10 K to be  $\sim 0.019 \pm 0.001$ .
- The derived values for the effective chemical desorption cross sections and efficiency per incident H strengthen the argument that H<sub>2</sub>S ice is effectively transferred to the gas phase through the excess energy generated by reactions with hydrogen atoms.
- The confirmation of H<sub>2</sub>S<sub>2</sub> formation as a result of HS radical recombination proves that non-energetic sulfur chemistry can take place at temperatures as low as 10 K through radical-radical reactions, which could potentially lead to the formation of sulfur-bearing COMs in more representative interstellar ice mixtures.
- We derived the effective formation cross sections for H<sub>2</sub>S<sub>2</sub> of  $(9.8 \pm 0.9) \times 10^{-17} \text{ cm}^2$ ,  $(7.8 \pm 0.9) \times 10^{-17} \text{ cm}^2$ ,  $(8.3 \pm 0.7) \times 10^{-17} \text{ cm}^2$ , and  $(5.2 \pm 0.6) \times 10^{-17} \text{ cm}^2$  at 10, 12, 14, and 16 K, respectively.
- No chemical desorption was observed upon formation of H<sub>2</sub>S<sub>2</sub> above the current detection limit.
- Approximately 85% to 74% of the H<sub>2</sub>S ice destruction observed under our experimental conditions can be associated with chemical desorption, whereas  $\sim 15\%$  to 26% is due to H<sub>2</sub>S<sub>2</sub> formation. The relative consumption of H<sub>2</sub>S by the latter process grows with temperature, implying that sulfur chemistry induced by HS radicals becomes increasingly more relevant in warmer environments.

*Acknowledgements.* This work has been supported by the Danish National Research Foundation through the Center of Excellence “InterCat” (Grant agreement no.: DNRF150); the Netherlands Research School for Astronomy (NOVA); and the Dutch Astrochemistry Network II (DANII). K.J.C. is grateful for support from NWO via a VENI fellowship (VI.Veni.212.296).

## References

- Bacmann, A., Taquet, V., Faure, A., Kahane, C., & Ceccarelli, C. 2012, *A&A*, **541**, A12
- Bertin, M., Romanzin, C., Doronin, M., et al. 2016, *ApJ*, **817**, L12
- Biver, N., Bockelée-Morvan, D., Moreno, R., et al. 2015, *Sci. Adv.*, **1**, 1500863
- Bull, J. N., & Harland, P. W. 2008, *Int. J. Mass Spectrom.*, **273**, 53
- Bull, J. N., Harland, P. W., & Vallance, C. 2012, *J. Phys. Chem. A*, **116**, 767
- Calmonte, U., Altwegg, K., Balsiger, H., et al. 2016, *MNRAS*, **462**, S253
- Cazaux, S., Cobut, V., Marseille, M., Spaans, M., & Caselli, P. 2010, *A&A*, **522**, A74
- Cazaux, S., Carrascosa, H., Muñoz Caro, G. M., et al. 2022, *A&A*, **657**, A100
- Cernicharo, J., Marcelino, N., Roueff, E., et al. 2012, *ApJ*, **759**, A43
- Chen, Y. J., Chuang, K. J., Muñoz Caro, G. M., et al. 2014, *ApJ*, **781**, 15
- Chen, Y. J., Juang, K. J., Nuevo, M., et al. 2015, *ApJ*, **798**, 80
- Chuang, K.-J. 2018, Ph.D. Thesis, University of Leiden, The Netherlands
- Chuang, K. J., Fedoseev, G., Qasim, D., et al. 2018, *ApJ*, **853**, 102
- Collings, M. P., Anderson, M. A., Chen, R., et al. 2004, *MNRAS*, **354**, 1133
- Cruz-Díaz, G. A., Martín-Doménech, R., Muñoz Caro, G. M., & Chen, Y. J. 2016, *A&A*, **592**, A68
- Cuppen, H., Ioppolo, S., Romanzin, C., & Linnartz, H. 2010, *Phys. Chem. Chem. Phys.*, **12**, 12077
- Cuppen, H. M., Walsh, C., Lamberts, T., et al. 2017, *Space Sci. Rev.*, **212**, 1
- Dulieu, F., Congiu, E., Noble, J., et al. 2013, *Sci. Rep.*, **3**, 1338
- Fayolle, E. C., Bertin, M., Romanzin, C., et al. 2011, *ApJ*, **739**, L36
- Fredon, A., Radchenko, A. K., & Cuppen, H. M. 2021, *Acc. Chem. Res.*, **54**, 745
- Fuchs, G. W., Cuppen, H. M., Ioppolo, S., et al. 2009, *A&A*, **505**, 629
- Furuya, K., Oba, Y., & Shimonishi, T. 2022, *ApJ*, **926**, 171
- Garozzo, M., Fulvio, D., Kanuchova, Z., Palumbo, M. E., & Strazzulla, G. 2010, *A&A*, **509**, A67
- Garrod, R., Park, I. H., Caselli, P., & Herbst, E. 2006, *Faraday Discuss.*, **133**, 51
- Garrod, R. T., Wakelam, V., & Herbst, E. 2007, *A&A*, **467**, 1103
- Hatchell, J., Thompson, M. A., Millar, T. J., & MacDonald, G. H. 1998, *A&A*, **338**, 713
- He, J., Emtiaz, S. M., & Vidali, G. 2017, *ApJ*, **851**, 104
- Hollenberg, J. L., & Dows, D. A. 1961, *J. Chem. Phys.*, **34**, 1061
- Hudson, J. E., Hamilton, M. L., Vallance, C., & Harland, P. W. 2003, *Phys. Chem. Chem. Phys.*, **5**, 3162



- Hudson, J. E., Weng, Z. F., Vallance, C., & Harland, P. W. 2006, *Int. J. Mass Spectrom.*, **248**, 42
- Ioppolo, S., Cuppen, H. M., Romanzin, C., van Dishoeck, E. F., & Linnartz, H. 2008, *ApJ*, **686**, 1474
- Ioppolo, S., Cuppen, H. M., Romanzin, C., van Dishoeck, E. F., & Linnartz, H. 2010, *Phys. Chem. Chem. Phys.*, **12**, 12065
- Ioppolo, S., Fedoseev, G., Lamberts, T., Romanzin, C., & Linnartz, H. 2013, *Rev. Sci. Instrum.*, **84**, 073112
- Isoniemi, E., Pettersson, M., Khriachtchev, L., Lundell, J., & Räsänen, M. 1999, *J. Phys. Chem. A*, **103**, 679
- Jiménez-Escobar, A., & Muñoz Caro, G. M. 2011, *A&A*, **536**, A91
- Jiménez-Escobar, A., Muñoz Caro, G. M., & Chen, Y. J. 2014, *MNRAS*, **443**, 343
- Jiménez-Serra, I., Vasyunin, A. I., Caselli, P., et al. 2016, *ApJ*, **830**, L6
- Laas, J. C., & Caselli, P. 2019, *A&A*, **624**, A108
- Lamberts, T., & Kästner, J. 2017, *J. Phys. Chem. A*, **121**, 9736
- Le Roy, L., Altwegg, K., Balsiger, H., et al. 2015, *A&A*, **583**, A1
- Martín-Doménech, R., Manzano-Santamaría, J., Muñoz Caro, G. M., et al. 2015, *A&A*, **584**, A14
- McClure, M. K., Rocha, W. R. M., Pontoppidan, K. M., et al. 2023, *Nat. Astron.*, **7**, 431
- McCord, T. B., Hansen, G. B., Clark, R. N., et al. 1998, *J. Geophys. Res.*, **103**, 8603
- Mífsud, D. V., Herczku, P., Rácz, R., et al. 2022, *Front. Chem.*, **10**, 1003163
- Minh, Y. C., Irvine, W. M., & Ziurys, L. M. 1989, *ApJ*, **345**, L63
- Minissale, M., & Dulieu, F. 2014, *J. Chem. Phys.*, **141**, 014304
- Minissale, M., Moudens, A., Baouche, S., Chaabouni, H., & Dulieu, F. 2016, *MNRAS*, **458**, 2953
- Moore, M., Hudson, R., & Carlson, R. 2007, *Icarus*, **189**, 409
- Muñoz Caro, G. M., Jiménez-Escobar, A., Martín-Gago, J. Á., et al. 2010, *A&A*, **522**, A108
- Nash, D. B., & Howell, R. R. 1989, *Science*, **244**, 454
- Navarro-Almaida, D., Le Gal, R., Fuente, A., et al. 2020, *A&A*, **637**, A39
- Neufeld, D. A., Godard, B., Gerin, M., et al. 2015, *A&A*, **577**, A49
- Nguyen, T., Oba, Y., Shimonishi, T., Kouchi, A., & Watanabe, N. 2020, *ApJ*, **898**, L52
- Nguyen, T., Oba, Y., Sameera, W. M. C., Kouchi, A., & Watanabe, N. 2021, *ApJ*, **918**, 73
- Oba, Y., Tomaru, T., Lamberts, T., Kouchi, A., & Watanabe, N. 2018, *Nat. Astron.*, **2**, 228
- Oba, Y., Tomaru, T., Kouchi, A., & Watanabe, N. 2019, *ApJ*, **874**, 124
- Öberg, K. I., Fuchs, G. W., Awad, Z., et al. 2007, *ApJ*, **662**, L23
- Öberg, K. I., Bottinelli, S., Jørgensen, J. K., & van Dishoeck, E. F. 2010, *ApJ*, **716**, 825
- Paardekooper, D. M., Fedoseev, G., Riedo, A., & Linnartz, H. 2016, *A&A*, **596**, A72
- Phuong, N. T., Chapillon, E., Majumdar, L., et al. 2018, *A&A*, **616**, A5
- Qasim, D., Witlox, M. J. A., Fedoseev, G., et al. 2020, *Rev. Sci. Instrum.*, **91**, 054501
- Santos, J. C., Chuang, K.-J., Schrauwen, J. G. M., et al. 2023, *A&A*, **672**, A112
- Scibelli, S., & Shirley, Y. 2020, *ApJ*, **891**, 73
- Shingledecker, C. N., Lamberts, T., Laas, J. C., et al. 2020, *ApJ*, **888**, 52
- Sie, N.-E., Cho, Y.-T., Huang, C.-H., et al. 2022, *ApJ*, **938**, 48
- Smith, R. G. 1991, *MNRAS*, **249**, 172
- Thaddeus, P., Kutner, M. L., Penzias, A. A., Wilson, R. W., & Jefferts, K. B. 1972, *ApJ*, **176**, L73
- Tielens, A. G. G. M., & Hagen, W. 1982, *A&A*, **114**, 245
- Tschersich, K. G. 2000, *J. Appl. Phys.*, **87**, 2565
- van der Tak, F. F. S., Boonman, A. M. S., Braakman, R., & van Dishoeck, E. F. 2003, *A&A*, **412**, 133
- van Dishoeck, E. F., Blake, G. A., Jansen, D. J., & Groesbeck, T. D. 1995, *ApJ*, **447**, 760
- van Dishoeck, E. F., Herbst, E., & Neufeld, D. A. 2013, *Chem. Rev.*, **113**, 9043
- Vastel, C., Phillips, T. G., Ceccarelli, C., & Pearson, J. 2003, *ApJ*, **593**, L97
- Vasyunin, A. I., & Herbst, E. 2013, *ApJ*, **769**, 34
- Vidal, T. H. G., Loison, J.-C., Jaziri, A. Y., et al. 2017, *MNRAS*, **469**, 435
- Wakelam, V., Castets, A., Ceccarelli, C., et al. 2004, *A&A*, **413**, 609
- Watanabe, N., & Kouchi, A. 2002, *ApJ*, **571**, L173
- Watanabe, N., & Kouchi, A. 2008, *Progr. Surf. Sci.*, **83**, 439
- Westley, M. S., Baratta, G. A., & Baragiola, R. A. 1998, *J. Chem. Phys.*, **108**, 3321
- Yarnall, Y. Y., & Hudson, R. L. 2022, *ApJ*, **931**, L4

## Appendix A: Determination of IR band strengths

The derivation of  $A'(\text{H}_2\text{S})$  for our specific experimental settings was performed in a manner similar to what was described by [Chuang et al. \(2018\)](#). Infrared spectra are acquired during deposition of an  $\text{H}_2\text{S}$  ice at 10 K while simultaneously measuring the interference pattern of a HeNe laser that hits the ice sample at an incident angle of  $\sim 1.7^\circ$ . The thickness of the ice ( $d_X$ ) is derived from the laser fringe pattern by the equation ([Hollenberg & Dows 1961](#); [Westley et al. 1998](#)):


$$d_X = k \times \frac{\lambda}{2n_X \cdot \cos(\theta_f)}, \quad (\text{A.1})$$

where  $k$  is the number of fringes,  $\lambda$  is the laser wavelength (i.e., 632.8 nm),  $n_X$  is the refractive index of the ice species, and  $\theta_f$  is the angle of refraction in degrees. From the thickness measurements, it is possible to derive the absolute column density of the ice by the equation:

$$N_X = \frac{d_X \cdot \rho_X \cdot N_a}{M_X}, \quad (\text{A.2})$$

where  $\rho_X$  is the density in grams per cubic centimeter,  $N_a$  is the Avogadro's constant, and  $M_X$  is the molar mass of a given species.

In the left panel of [Fig. A.1](#), the HeNe laser interference pattern is shown as a function of time and fit with a sinusoidal function. This pattern arises from the growing  $\text{H}_2\text{S}$  ice being deposited on the sample. The corresponding increase in the IR absorption area of the SH stretching mode of  $\text{H}_2\text{S}$  (i.e.,  $\sim 2553 \text{ cm}^{-1}$ ) as a function of ice column density is shown in the right panel of [Fig. A.1](#). A linear fit to the points yields a band strength value in reflection mode and is specific to our experimental setup of  $A'(\text{H}_2\text{S})_{\sim 2553 \text{ cm}^{-1}} \sim (4.7 \pm 0.1) \times 10^{-17} \text{ cm molecule}^{-1}$ . For this calculation, we utilized  $\rho = 0.944 \pm 0.005 \text{ g cm}^{-3}$  and  $n = 1.407 \pm 0.005$ , as reported by [Yarnall & Hudson \(2022\)](#).



**Fig. A.1.** Derivation of  $A'(\text{H}_2\text{S})$ . Left panel: HeNe laser interference pattern obtained from the growing  $\text{H}_2\text{S}$  ice at 10 K (black) together with its sinusoidal fitting (red). Changes in the amplitude of the fringes are attributed to laser instabilities. Right panel: Integrated absorbance area of the SH stretching mode of  $\text{H}_2\text{S}$  as a function of column density as derived from the fringe pattern. A linear fit to the points is also shown in red.